



Head Start to A-Level Maths

Bridge the gap between GCSE and A-Level



CGP is the <u>only</u> choice for A-Level Maths!



Unbeatable books and practice papers for A-Level Maths and Further Maths — revision essentials for **AQA**, **Edexcel**, **OCR** and **OCR MEI**

cgpbooks.co.uk



Head Start to A-Level Maths

Looking to start your A-Level Maths studies early? Well, let's get this show on the road with this unbeatable CGP Head Start book!

Step 1: Recap key GCSE concepts (in case you've already forgotten!). Step 2: Introduce new A-Level ideas with **clear study notes** and **practice questions**. Step 3: Strut into A-Level Maths with confidence!

CGP — still the best! ③

Our sole aim here at CGP is to produce the highest quality books — carefully written, immaculately presented and dangerously close to being funny.

Then we work our socks off to get them out to you — at the cheapest possible prices.

Contents

Section 1 — Number

Types of Number	6
Fractions	7

Section 2 — Basic Algebra

Laws of Indices	
Multiplying Out Brackets	
Factorising	
Surds	
Solving Equations	
Rearranging Formulas	

Section 3 — Quadratic Equations

Factorising Quadratics	.16
The Quadratic Formula	.18
Completing the Square	.19

Section 4 — More Algebra

lgebraic Fractions	.22
nequalities	.24
Graphical Inequalities	.26
imultaneous Equations	.28
roof	.30
unctions	.32

Section 5 — Graphs

Straight Lines	34
Parallel and Perpendicular Lines	
Quadratic Graphs	37
Harder Graphs	
Graph Transformations	40

Section 6 — Trigonometry and Vectors

Trigonometry — Sin, Cos, Tan	42
Trigonometry — Graphs	44
The Sine and Cosine Rules	47
Vectors	49

Section 7 — Statistics and Probability

Sampling	51
Data Basics	
Histograms	53
Averages	54
Cumulative Frequency	
Probability	57
Laws of Probability	
Tree Diagrams	60

nswers	61
ndex	76

Throughout the book, we've marked up content that you're not likely to have come across at GCSE with stamps like this one. This is the material you'll be learning more about as part of A-level Maths.



Published by CGP

Typesetters: Callum Lamb, Kirsty Sweetman

Editors: Rachel Craig-McFeely, Sarah Pattison

MBR72 \sim 0521 $\scriptstyle \sim$ 22597

With thanks to Emma Clayton for the proofreading. With thanks to Emily Smith for the copyright research.

Clipart from Corel®

Based on the classic CGP style created by Richard Parsons.

Text, design, layout and original illustrations © Coordination Group Publications Ltd. (CGP) 2021 All rights reserved.

Extra copies are available from CGP with next day delivery • 0800 1712 712 • www.cgpbooks.co.uk

This first exercise will help you find out which areas of maths you need to work on before you start your A-Level Maths course. Do it before you work through the book — if you struggle with any of the questions, go straight to the relevant pages to brush up on your skills.

Diagnostic Test

Once you've done all that, work through the rest of the book. You'll be able to recap and practise some useful GCSE topics and see how they'll lead into your A-Level work.

Ту	pes of Number and Fractions	These topics a	re covered in Section	1 — p.6-7. =
1)	Which of the following are integers?			
	4 -3.5 0.3 $\frac{4}{5}$	8.99	-10	205 0
2)	Which of the following values are rational, ar	nd which are irration	nal?	
	5.9 $\pi \sqrt{7} \frac{1}{5}$	-6 $\sqrt{4}$	13.978	2.1
3)	Evaluate the following without using a calcul Give any answers larger than 1 as improper f	ator, giving your ans ractions.	wers in their lowe	est terms.
	a) $\frac{2}{9} \times \frac{3}{5}$ b) $\frac{1}{6} \div \frac{2}{3}$	c) $\frac{1}{12} + \frac{5}{6}$	d) $\frac{8}{5}$ –	$\frac{1}{7}$
Inc	lices, Multiplying Out Brackets	and Factorisi	ing	e topics are
4)	Simplify the following:a) $x^7 \times x^2$ b) $10y^3 \div 5y$	c) <i>m</i> ⁰		d) $(2n^2)^5$
5)	Write 5^{-2} as a fraction.			
6)	Evaluate the following without using a calcul	ator:		
	a) $\left(\frac{3}{4}\right)^2$ b) $16^{\frac{1}{2}}$	c) $8^{\frac{2}{3}}$		d) $36^{-\frac{1}{2}}$
7)	Multiply out the brackets and simplify your as a) $(x + 4)(x - 6)$ b) $(x + 5)^2$	nswers where possib c) $(2x - 1)(x + 3)$	d) (x +	1)(x-4)(x+5)
8)	Factorise the following:	-) 2 4		-) 2 F
	a) $5x + 20$ b) $3a + 12ab$	C) $x^2 - 4$	d) $9x^2 - 36$	e) $x^2 - 5$
Su	ras 7/11/11/11/11/11/11/11/11/11/11/11/11/11			
9)	Simplify the following:	$\sqrt{20}$		
	a) $\sqrt{3} \times \sqrt{2}$ b) $(\sqrt{5})^2$ c)	$\frac{\sqrt{50}}{\sqrt{6}}$ d)	$\sqrt{12} + 2\sqrt{3}$	e) $(1 + \sqrt{7})^2$
10)	Rationalise the denominators of the following	<u>.</u>		-
	a) $\frac{3}{\sqrt{2}}$ b) $\frac{\sqrt{5}}{2\sqrt{2}}$	c) $\frac{2}{3 + \sqrt{6}}$	d) $\frac{\sqrt{2}}{1-\sqrt{2}}$	<u>-</u>

TEST YOURSELF

Diagnostic Test

TEST YOURSELF

Diagnostic Test



2

Diagnostic Test

TEST YOURSELF

Proof and Functions

These topics are covered on p.30-33. =

- 22) Prove that the sum of any three consecutive odd numbers is a multiple of 3.
- 23) Naveen says, "for any integers x and y, xy > y". Prove that Naveen is wrong.
- 24) $f(x) = \frac{x+5}{3}$ and g(x) = x 3. a) Evaluate f(4).

b) Find fg(x).

c) Find $f^{-1}(x)$.

Straight Lines and Quadratic Graphs

- 25) Give the gradient and *y*-intercept of the line x + 2y = 4.
- 26) Point A has coordinates (5, 2) and point B has coordinates (2, -4).a) Find the equation of the line passing through points A and B.b) Find the exact length of line AB.
- Go to p.34-37 if you found = these questions tricky.

- 27) Line A has equation y = 2x + 5.
 - a) Find the equation of the line parallel to line A which passes through (3, 2).
 - b) Find the equation of the line perpendicular to line A which passes through (2, 1).
- 28) Sketch the graph of $y = x^2 8x + 15$. Label the graph with the coordinates of the turning point and the points where the graph crosses the axes.

Harder Graphs and Graph Transformations

29) Sketch the following graphs:

a)
$$y = x^3$$
 b) $y = \frac{1}{x}$ c) $y = -\frac{1}{x}$

- 30) The graph on the right shows how the number of fish (*F*) living in a river changes over time. The equation of the graph is $F = mn^t$ where *t* is the number of years and *m* and *n* are positive constants. Find the values of *m* and *n*.
- 31) Find the equation of the tangent to $x^2 + y^2 = 25$ at the point (3, 4). Give your answer in the form ax + by + c = 0.





32) $f(x) = x^2$. For parts a) to c) below, sketch the graphs of y = f(x) and the given transformation. a) y = f(x) + 3 b) y = f(x + 3) c) y = -f(x)

Diagnostic Test

Trigonometry and Vectors

TEST YOURSELF

These topics are in Section 6 — p.42-50. =

33) Find the unknowns in each of these triangles. Give your answers to 1 decimal place.



34) Find the unknowns in each of these triangles. Give your answers to 1 decimal place.



35) ABCD is the parallelogram shown on the right. M, N, P and Q are the midpoints of the sides. $\overline{AB} = \mathbf{a}$ and $\overline{BC} = \mathbf{b}$.

Find the following vectors in terms of **a** and **b**.

- a) \overrightarrow{AC} b) \overrightarrow{DQ} c) \overrightarrow{CM} d) \overrightarrow{OP} e) \overrightarrow{MB} f) \overrightarrow{PA}
- 36) The diagram shows triangle ABC. M is the midpoint of \overrightarrow{AC} and N is the midpoint of \overrightarrow{BC} . $\overrightarrow{AM} = 3\mathbf{a} - \mathbf{b}$ and $\overrightarrow{NC} = 2(\mathbf{a} - \mathbf{b})$. Show that \overrightarrow{AB} and \overrightarrow{MN} are parallel.







Sampling and Histograms

- 37) Describe how a simple random sample of size 20 can be selected from a population of 200.
- 38) The weights of the chocolate bars in a shop storeroom are shown in the table and histogram below.
 - a) Use the information in the table and the histogram to label the vertical axis.
 - b) Use the histogram to complete the table.
 - c) Use the table to add the missing bar to the histogram.

Weight (<i>w</i> , in grams)	Frequency
$0 < w \le 100$	50
$100 < w \le 150$	100
$150 < w \le 200$	150
$200 < w \le 250$	



Diagnostic Test

1

2

Averages and Cumulative Frequency

39) Find the mean, median and mode(s) of these numbers:

- 40) The table shows the journey times between home and school for 60 students.
 - a) Write down the modal class.
 - b) Which group contains the median?
 - c) Estimate the mean value.

5

- d) Draw a cumulative frequency graph for the data in the table.
- 41) Using this cumulative frequency graph, find the:



200000000000000000000000000000000000000
- Averages are covered on n54-55 -
- Twelages are covered on p.51.55.

4 2 6 11 -4

Time (<i>m</i> minutes)	Frequency
$5 < m \le 10$	4
$10 < m \le 15$	25
$15 < m \le 20$	18
$20 < m \leq 25$	8
$25 < m \leq 30$	5

a) median

b) lower quartile

1

- c) upper quartile
- d) interquartile range

You can learn about cumulative frequency on p.56.

Probability and Tree Diagrams



<u>NUTURE CONTRACTOR CONTRA TOR CONTRA TOR CON</u>

E You can learn about these topics on p.57-60. =

42) Lewis asked 50 people if they like mashed potatoes (M) and roast potatoes (R). The Venn diagram shows the results.

A person is chosen at random.

Find the probability that they:

- a) like mashed potatoes
- b) like neither mashed nor roast potatoes
- c) like both types of potatoes
- d) don't like roast potatoes
- e) don't like mashed potatoes
- 43) Mona's purse contains two £5 notes, four £10 notes and three £20 notes. It also contains five 20p coins, four 50p coins and three £1 coins.
 - a) Mona picks one note and one coin at random from her purse. Find the probability that she picks a £5 note and a 20p coin.
 - b) Mona picks two coins at random without replacement. Use a tree diagram to find the probability she picks a 50p coin and a £1 coin.

5

Types of Number

First things first, here are some **important terms** that will crop up **throughout A-Level Maths**. If you're clear on what they mean now, it'll make A-Level life a lot easier.

Integers are just **Whole Numbers**

- An **integer** is any positive or negative **whole number** (including **zero**). 1)
- 2) The set of integers is represented by the symbol \mathbb{Z} .

Rational Numbers can be Written as Fractions

- 1) A rational number is any number which can be written as a fraction with integers on the top and bottom. Don't forget, any integer can be written as a fraction over 1.
- 2) The set of **rational numbers** is represented by the symbol \mathbb{Q}
- 3) Both **terminating** and **recurring decimals** are rational.
- 4) A number which **cannot** be written exactly as a fraction is **irrational**. Irrational numbers are **non-repeating** decimals, which **never end**.
- For example, π is a **never-ending**, **non-repeating** decimal (π = 3.1415926535...). 5) This means that π is **irrational**.

EXAMPLES

Is 0.6 a rational or irrational number?

0.6 is the recurring decimal 0.6666666... Recurring decimals are rational $-0.6 = \frac{2}{3}$.

Is 5.26 a rational or irrational number?

5.26 is **rational** because it is a terminating decimal $\left(5.26 = \frac{526}{100} = \frac{263}{50}\right)$

All Rational and Irrational Numbers are Real Numbers

- 1) Any rational or irrational number is a real number.
- 2) The set of **real numbers** is represented by the symbol \mathbb{R} .
- 3) There are also numbers called **imaginary numbers.** They're **not real numbers**.
- Imaginary numbers are the result of taking the square root of a negative number. 4) It's not possible to do this with real numbers because the square of a real number is always positive, so **imaginary numbers** are used in these cases. You'll learn about them if you do A-Level Further Maths.

Rational? Irrational? Whatever man — I just keep it real...

- 1) Are these numbers rational or irrational? Explain your answer.
- b) 0.1478 c) $\sqrt{64}$ d) √3 a) 0.236849
- e) 2π 2) True or false? a) All integers are rational. b) A recurring decimal is irrational.



Alvin swore he wasn't asleep, he

was just thinking about integers.



Fractions

Multiplying Fractions — just Multiply the Numbers

- 1) Multiply the numerators and denominators separately.
- 2) Make sure to turn any **mixed numbers** into **improper fractions** first.
- 3) Cancel the fractions down before you start multiplying if you can it'll make things easier.

EXAMPLE

Find $1\frac{3}{4} \times 2\frac{2}{3}$.

Write the mixed numbers as improper fractions: Now cancel down across your fractions and multiply.

8 divides by 4, so you can cancel these down.

Dividing by a Fraction — **Flip It** and **Multiply**

- To divide by a fraction, just turn the fraction you're dividing by upside down, and change the divide sign to a multiply.
- 2) Just like before, turn **mixed numbers** into **improper fractions** and do any **cancelling** before you start.

 $1\frac{3}{4} \times 2\frac{2}{3} = \frac{4+3}{4} \times \frac{6+2}{3} = \frac{7}{4} \times \frac{8}{3}$ $\frac{7}{4} \times \frac{8}{3} = \frac{7}{1} \times \frac{2}{3} = \frac{14}{3}$

EXAMPLE Calculate $\frac{3}{8} \div \frac{7}{16}$.

> Flip the second fraction and multiply. Make sure you do any cancelling you can before multiplying. 16 divides by 8, so you can cancel these down: $\frac{3}{8} \div \frac{7}{16} = \frac{3}{8_1} \times \frac{16^2}{7} = \frac{3}{1} \times \frac{2}{7} = \frac{6}{7}$

The **Denominators** need to **Match** for **Addition** and **Subtraction**

- 1) In order to **add** or **subtract** fractions, the **denominators** must be the **same**. So you have to find the **lowest common multiple** of all **denominators** (the lowest common denominator).
- 2) You still need to turn any mixed numbers into improper fractions.
- 3) Once both fractions have a common denominator, you can add/subtract their numerators.

EXAMPLE

Find $1\frac{5}{6} - \frac{3}{4}$.

You'll need to be confident with all of these rules as they're really important for using algebraic fractions.

Convert the mixed number to an improper fraction: $1\frac{5}{6} - \frac{3}{4} = \frac{11}{6} - \frac{3}{4}$. Now they need a common denominator. The LCM of 6 and 4 is 12, so $\frac{11}{6} - \frac{3}{4} = \frac{22}{12} - \frac{9}{12}$. This means you can now subtract the numerators: $\frac{22}{12} - \frac{9}{12} = \frac{13}{12}$.



Laws of Indices

Indices are Powers

For the value 4³, 4 is the **base** and 3 is the **index**. (The plural of index is **indices**.) You'll use indices all the time in A-Level Maths — especially for topics like **algebra**, **differentiation** and **integration**. So make sure you know the **laws of indices** inside out...

Multiplying Indices = Add the Powers, Dividing = Subtract them

So $x^a \times x^b = x^{(a+b)}$ and $x^a \div x^b = x^{(a-b)}$

EXAMPLES

Simplify $a^4 \times a^2$.

Just add the indices: $a^4 \times a^2 = a^{(4+2)} = a^6$. You can see how this works by rewriting the calculation as one big multiplication: $a^4 \times a^2 = (a \times a \times a \times a) \times (a \times a) = a \times a \times a \times a \times a \times a = a^6$

Rewrite 45 ÷ 43 as a single power of 4.

This time, subtract the indices: $4^5 \div 4^3 = 4^{(5-3)} = 4^2$ Again, you can see how this works by rewriting the calculation: $4^5 \div 4^3 = \frac{4 \times 4 \times 4 \times 4}{4 \times 4 \times 4} = 4 \times 4 = 4^2$

Remember — this law only works when the two values have the same base. So $2^4 \times 2^7 = 2^{(4+7)} = 2^{11}$ is fine, but you can't use this rule to work out $2^4 \times 3^5$.

Simplify $8a^5 \times 2a^6$.

Both terms have the same base, *a*, and multiplication is **commutative** (which means it doesn't matter what order you do it in). So you can rewrite this as: $8 \times a^5 \times 2 \times a^6 = (8 \times 2) \times (a^5 \times a^6) = 16 \times a^{(5+6)} = 16a^{11}$

Simplify $(x - 1)^9 \div (x - 1)^4 \times y^3 \times y^4$.

Deal with the powers of each base separately: $(x - 1)^9 \div (x - 1)^4 \times y^3 \times y^4 = (x - 1)^{(9-4)} \times y^{(3+4)} = (x - 1)^5 y^7$ Don't be put off by the brackets (x - 1) is just the base.

There are **Rules** for **x⁰** and **x¹**

 $x^0 = 1$ for any value of x The easiest way to see this is this with an example: $4^3 \div 4^3 = 4^{(3-3)} = 4^0$ and $4^3 \div 4^3 = 1$, so $4^0 = 1$.

 $x^1 = x$ for any value of x Again, this is easiest to understand as an example: $4^3 \div 4^2 = 4^{(3-2)} = 4^1$, and $4^3 \div 4^2 = 64 \div 16 = 4$, so $4^1 = 4$.

Revise more I tell you — sorry these powers have gone to my head...

1) Simplify the following:

I)	Simplify the to	
	a) $b^5 \times b^6$	
	e) $x^2 \times x^3 \div x^4$	

b) $a^9 \times a \times b^5$ f) $z^3 \times (y+2)^5 \times z \div (y+2)^2$

c) $c^5 \div c^2$ g) $a^{(x+2)} \times a^{2x}$

Laws of Indices

To Raise a Power to Another Power, Multiply them

So $(x^a)^b = x^{ab}$

EXAMPLES Simplify $(q^3)^2$. $(q^3)^2 = q^{(3 \times 2)} = q^6$

You can see how this works by	
= rewriting it as one big multiplication:	
$= (q^3)^2 = q^3 \times q^3 = (q \times q \times q) \times (q \times q \times q) =$	
$= q \times q \times q \times q \times q \times q = q^{6}$	

For a fraction raised to a negative power, you turn the fraction upside down, then apply the positive index.

Express 26 ÷ 42 as a single power.

You need to make the bases the same before simplifying, so rewrite 4 as a power of 2 using the rule above: $2^6 \div 4^2 = 2^6 \div (2^2)^2 = 2^6 \div 2^{(2 \times 2)} = 2^6 \div 2^4 = 2^2$

Write $\frac{1}{125}$ as a power of 5.

 $125 = 5^3$, so $\frac{1}{125} = \frac{1}{5^3} = 5^{-3}$

Negative and **Fractional Indices** are a bit **Trickier**

A negative index means '1 ÷ the positive power' — so $x^{-a} = \frac{1}{x^{a}}$

EXAMPLES

Write 3⁻⁴ as a fraction.

$$3^{-4} = \frac{1}{3^4} = \frac{1}{81}$$

If a number has a **fractional** index, this means '**the root of**' — so $x^{\frac{1}{a}} = \sqrt[a]{x}$

EXAMPLES

Find $64^{\frac{1}{3}}$ without using a calculator.

 $64^{\frac{1}{3}} = \sqrt[3]{64}$ This is the cube root of 64, so the answer is **4**. (Because $4 \times 4 \times 4 = 64$.)

Find $\left(\frac{1}{256}\right)^{\frac{1}{4}}$ without using a calculator.

When you raise a fraction to a power, you raise the top and bottom to that power:

$$\left(\frac{1}{256}\right)^{\frac{1}{4}} = \frac{1^{\frac{1}{4}}}{256^{\frac{1}{4}}} = \frac{4/1}{4\sqrt{256}} = \frac{1}{4}$$

You might have to **rewrite** the index. Use the fact that $x^{\frac{a}{b}} = (x^{\frac{1}{b}})^a$, which is the same as $(x^a)^{\frac{1}{b}}$.

If you get a negative fractional index, use this fact to turn $x^{-\frac{a}{b}}$ into $\left(x^{\frac{a}{b}}\right)^{-1}$ or $(x^{-1})^{\frac{a}{b}}$.



Find $27^{-\frac{1}{3}}$ without using a calculator.

$$27^{-\frac{1}{3}} = (27^{\frac{1}{3}})^{-1} = (\sqrt[3]{27})^{-1} = (3)^{-1} = \frac{1}{3}$$



Multiplying Out Brackets

Double Brackets and Squared Brackets Can Be Expanded

When expanding double brackets, multiply each term in one set 1) of brackets by each term in the other. If each set of brackets contains two terms, you can use the FOIL method to make sure you don't miss any terms:

Multiply the **First** term in each set of brackets together, then the two Outside terms, then the two Inside terms and finally the Last terms.

2) Write out squared brackets as double brackets and then use the FOIL method to avoid making silly mistakes.

EXAMPLE

Expand and simplify $(2x - 4)^2$.

 $(2x - 4)^2 = (2x - 4)(2x - 4) = (2x \times 2x) + (2x \times -4) + (-4 \times 2x) + (-4 \times -4)$ $= 4x^2 - 8x - 8x + 16 = 4x^2 - 16x + 16$

Use a Similar Method with Triple Brackets

- 1) You can think of **triple brackets** as **double brackets** multiplied by **one more** set of brackets.
- So, use the **double bracket method** on two of the sets, 2) then multiply the **result** by the **remaining set of brackets**.
- You'll probably end up with more than two terms in your 'result', so you won't be able 3) to use the FOIL method for the last bit. Instead, you should **break up** the **shorter** set of brackets so that each of its terms is multiplied by the **second** set of brackets **separately**.
- 4) Then multiply out each of the new sets of brackets **one at a time**.

EXAMPLE

Write (x + 5)(x - 1)(x + 2) as a cubic expression.

 $(x + 5)(x - 1)(x + 2) = (x + 5)(x^{2} + x - 2) = x(x^{2} + x - 2) + 5(x^{2} + x - 2)$ $= (x^{3} + x^{2} - 2x) + (5x^{2} + 5x - 10) = x^{3} + 6x^{2} + 3x - 10$

Brackets can have More Than Two Terms



At A-Level, you might be asked to multiply out sets of brackets with more than two terms. As usual, you need to multiply every term in the first set of brackets by every term in the second. The method for this is just the same as **stage two** of the **triple bracket method** above – multiply each term in the shorter set of brackets by the other set separately.

EXAMPLE

Expand and simplify the expression $(x^2 - 5x - 1)(x + y + 1)$.

 $(x^{2} - 5x - 1)(x + y + 1) = x^{2}(x + y + 1) + (-5x)(x + y + 1) + (-1)(x + y + 1)$ $= (x^{3} + x^{2}y + x^{2}) + (-5x^{2} - 5xy - 5x) + (-x - y - 1)$ $= x^{3} + x^{2}y - 4x^{2} - 5xy - 6x - y - 1$



Quintuple brackets — just pack it all in and go home...

Multiply out each of these sets of brackets: a) (y + 3)(y - 6)b) (a-3)(b+4)

c) $(p-1)(p-2q)^2$





These techniques are wells

These techniques are really

useful for lots of A-Level

topics — like suras and various types of equations.

S Don't make the milet of the second

Don't make the mistake of just

squaring the stuff inside the brackets. In general, $(a + b)^2 \neq a^2 + b^2$. $= \ln \text{ general, } (a + b)^2 \neq a^2 + b^2. = \frac{1}{2}$

> Whongar ...

Whenever you multiply

out brackets, make sure

you simplify at the end — the x terms combine here.

Tina had mixed up 'cubic expressions' with 'cubist expressionism'.

Factorising

Factorising is Putting Brackets Into an Expression

Factorising means finding **common factors** that are in each term in an expression. You can take common factors outside a set of brackets to **rewrite** an expression. It's a handy skill that's used all the time in A-Level Maths, e.g. in **algebraic fractions** and **solving quadratics** and **cubics**.

- 1) You have to spot **common factors** in each term. These then come out to the **front** of the expression, and you put what's left in **brackets**.
- 2) Make sure that after you've finished factorising, any sets of brackets **can't be factorised** any more.

EXAMPLES

```
Factorise 14x + 21xy.
```

14x + 21xy = 7x(2 + 3y)

Factorise $16x^3 + 4x^2 - 4x$.	
$16x^3 + 4x^2 - 4x = 4x(4x^2 + x - 1)$)

Check your answer by multiplying the brackets out again — if it's right, you'll end up back where you started...

At A-Level, you might see expressions where the common factor is a set of brackets.

EXAMPLE

Factorise 3x(x + 2) - 4(x + 2).

(x + 2) appears in both terms of the expression, so it's a common factor: 3x(x + 2) - 4(x + 2) = (x + 2)(3x - 4)

The Difference Of Two Squares is One Square Minus Another

Any expression of the form $a^2 - b^2$ is called the **difference of two squares**. It can be **factorised** really easily using this result:

$$a^2 - b^2 = (a + b)(a - b)$$

You can see why this works by multiplying out (a + b)(a - b): $(a + b)(a - b) = a^2 - ab + ba - b^2 = a^2 - ab + ab - b^2 = a^2 - b^2$.

EXAMPLES

Factorise $x^2 - 25$.

a = x and b = 5, so $x^2 - 25 = (x + 5)(x - 5)$

Factorise $4x^2 - 81$. $4x^2 = (2x)^2$, so $4x^2 - 81 = (2x + 9)(2x - 9)$

and cancel down algebraic fractions, so you need = to be able to spot when you can use it.

If the number isn't a square, you can write it as a square root squared (see the next page).

EXAMPLE

Factorise $x^2 - 3y^2$.

 $3y^2 = (\sqrt{3}y)^2$, so using the difference of two squares: $x^2 - 3y^2 = (x + \sqrt{3}y)(x - \sqrt{3}y)$.

X-Factorising — warbling live on telly to earn the public's affection... 1) Factorise the following expressions: a) $20x^2 - 4x$ b) $8x^2y + 28xy^2$ c) $3\pi a^2 + 4\pi ab + 2\pi a$ d) $5x^2(x-1) - 2x(x-1)$ e) $x^2 - 9$ f) $9x^2 - 25$ g) $p^2 - 49q^2$ h) $v^2 - 7u^2$

Surds

Surds are the Square Roots of Non-Square Numbers

Irrational numbers that can be written as **roots** ($\sqrt{}$) are called **surds**. You need to be able to deal with them when working with **quadratics**, **cubics** and **vectors**.



Surds in the Denominator should be Rationalised

If there's a surd on the **bottom** of a fraction, you need to **get rid** of it. To do this, you multiply the **top** and **bottom** of the fraction by an expression that will give a **rational number** in the denominator. This is called **rationalising the denominator**.

EXAMPLES

Rationalise the denominator of $\frac{2/2}{\sqrt{2}}$.

You need to multiply top and bottom by something that will make the denominator rational — $\sqrt{5}$ will work because $\sqrt{a}\sqrt{a} = a$:

$$\frac{2\sqrt{2}}{\sqrt{5}} = \frac{2\sqrt{2}\sqrt{5}}{\sqrt{5}\sqrt{5}} = \frac{2\sqrt{10}}{5}$$

Show that $\sqrt{80} + \frac{25}{\sqrt{5}} = 9\sqrt{5}$. $\sqrt{80} = \sqrt{16}\sqrt{5} = 4\sqrt{5}$ and $\frac{25}{\sqrt{5}} = \frac{25\sqrt{5}}{\sqrt{5}\sqrt{5}} = \frac{25\sqrt{5}}{5} = 5\sqrt{5}$, so: $\sqrt{80} + \frac{25}{\sqrt{5}} = 4\sqrt{5} + 5\sqrt{5} = 9\sqrt{5}$



Surds

You can Rationalise more Complicated Denominators

You'll need a slightly different method for rationalising more complicated denominators.

- 1) Just like before, you multiply the **top** and **bottom** of the fraction by the **same** expression.
- 2) The expression you need to multiply by is just the **denominator** with the **opposite sign** in front of the **surd**.

EXAMPLE

Rationalise the denominator of $\frac{3}{1+\sqrt{2}}$.

The bit you want to rationalise is $1 + \sqrt{2}$, so multiply the top and bottom by $1 - \sqrt{2}$:

$$\frac{3}{1+\sqrt{2}} = \frac{3(1-\sqrt{2})}{(1+\sqrt{2})(1-\sqrt{2})} = \frac{3-3\sqrt{2}}{1^2-(\sqrt{2})^2}$$

$$= \frac{3-3\sqrt{2}}{-1} = 3\sqrt{2} - 3$$
This comes from the difference of two squares rule in the previous page: $(a+\sqrt{b})(a-\sqrt{b}) = a^2 - b$, with $a = 1$ and $b = 2$. You could also just multiply the brackets out.

This trick also works when there's a **number in front** of the **root** in the denominator. Just choose an expression with the **opposite sign** in front of the number multiplied by the surd.

EXAMPLE

Rationalise the denominator of $\frac{3+\sqrt{5}}{3+2\sqrt{5}}$.

Although this looks harder, just do the same thing. The denominator is $3 + 2\sqrt{5}$, so multiply the top and bottom by $3 - 2\sqrt{5}$:

$$\frac{3+\sqrt{5}}{3+2\sqrt{5}} = \frac{(3+\sqrt{5})(3-2\sqrt{5})}{(3+2\sqrt{5})(3-2\sqrt{5})}$$

$$= \frac{3^2-6\sqrt{5}+3\sqrt{5}-2(\sqrt{5})^2}{3^2-6\sqrt{5}+6\sqrt{5}-(2\sqrt{5})^2}$$

$$= \frac{9-3\sqrt{5}-2\times5}{9-2^2\times5}$$

$$= \frac{-3\sqrt{5}-1}{-11} = \frac{3\sqrt{5}+1}{11}$$



Obasi was struggling to rationalise his fashion choices.



Use the 6-Step Method to Solve Equations

Lots of A-Level Maths requires you to be able to **solve equations**. The basic skills covered here crop up loads in **differentiation**, **trigonometry**, and solving **quadratics** and **cubics**. To solve an equation, follow the **steps** below — you can **ignore** any steps you don't need. This method works for **any variable** — but if you're solving for *x*:

- 1) Remove any **fractions** by multiplying by the denominator(s).
- 2) Multiply out any brackets (see page 10).
- 3) **Collect** all the *x*-terms on one side and all the number terms on the other.
- 4) By **combining like terms**, **reduce** it to the form 'Ax = B'.
- 5) Finally, **divide** both sides by **A**. This gives your answer 'x = '.
- 6) If you had ' x^2 = ' instead, **square root** both sides to end up with ' $x = \pm$ '.

EXAMPLE

Solve $\frac{2x}{3} + \frac{4-x}{4} = 7x$.

- 1) First, get rid of the fractions. Multiply by 3 and then 4 to cancel the denominators: $\frac{3 \times 4 \times 2x}{3} + \frac{3 \times 4 \times (4 - x)}{4} = 3 \times 4 \times 7x$, so 8x + 3(4 - x) = 84x
- 2) Next, multiply out the set of brackets: 8x + 12 3x = 84x
- 3) Now collect the *x*-terms on one side of the equation and the number terms on the other: 12 = 84x - 8x + 3x
- 4) Combine like terms: 79x = 12
- 5) And divide by 79 to get $x = \frac{12}{79}$
- You don't need step 6 because there's no x^2 term...

Here are a couple more examples — both have a **squared** term to deal with, so you'll need to use **step 6** of the method above.

Remember, when you take the **square root** of a number the answer can be **positive** or **negative**, so the equation will have **two solutions**.

EXAMPLES

Solve 3x(x + 4) = 12x + 6.

Multiply out the brackets: $3x^2 + 12x = 12x + 6$

Simplify: $3x^2 + 12x - 12x = 6$ $\Rightarrow 3x^2 = 6$

Divide by 3: $x^2 = 2$

Take the square root: $x = \pm \sqrt{2}$

Be careful — the context of a question might = mean only one of these answers will make sense. =

Solve $5a = \frac{125}{a}$.

Multiply by *a* to get rid of the fraction: $5a^2 = 125$

Divide by 5: $a^2 = 25$

So,
$$a = \pm \sqrt{25} = \pm 5$$

PRACTICE QUESTIONS

What does a square Dalek say? x²-term-inate*...

1) Solve each of the following equations:

a) 4(2x-3) = 7x b) 3(x+14) = x + 12 c) $\frac{b-7}{3} + \frac{b+1}{5} = -1$

d)
$$\frac{q(q+7)}{7} - q$$

2) The sides of a square are *x* cm long. The area of $\frac{1}{4}$ of the square is 25 cm². How long are the sides?

*I'm so sorry.

 $= 4 - q^2$

Rearranging Formulas

The **Subject** of a **Formula** is the Letter on its **Own**

- If a formula has a **single letter** on one side of the equals sign, this letter is known as 1) the **subject** of the formula. For example, *y* is the subject of the formula y = mx + c.
- Sometimes you'll need to rearrange a formula to make a different letter the subject. 2)
- 3) You'll use this a lot in the **mechanics** sections of A-Level Maths, especially with **constant acceleration formulas**, so make sure you've got it learned.

This method is the same as the one = for solving equations — so vouve Use These Handy 7 Steps to Rearrange Formulas

- Square both sides to get rid of any square roots. 1)
- 2) **Multiply** by the denominator(s) to get rid of any **fractions**.
- 3) Multiply out any brackets.
- 4) **Collect** all the **subject** terms together on one side and all **non-subject terms** on the other.
- 5) By **combining** like terms, **reduce** it to the form 'Ax = B'. You might have to do some **factorising** to get it in the right form.
- 6) **Divide** both sides by A to give 'x = '.
- 7) If you've got ' $x^2 = '$, square root both sides to get ' $x = \pm '$.

EXAMPLE

Make *n* the subject of the formula $\sqrt{\frac{n+3}{a}} = 2m$.

First, square both sides to get rid of the square root:

Then multiply by *a* to get rid of the fraction:

Finally, collect all the terms without an *n* on the RHS by subtracting 3: $n = 4am^2 - 3$

If the subject appears more than once in the formula, then you'll need to do some factorising.

 $\frac{n+3}{a} = 4m^2$

 $n + 3 = 4am^2$

Loyal subjects I command thee — moveth to yonder side of the '='...

EXAMPLE

Make *a* the subject of the formula $a^2 + b^2 = (3 + a)(a - b)$.

There aren't any square roots or fractions, so go straight to multiplying out the brackets: $a^2 + b^2 = 3a - 3b + a^2 - ab$.

Gather all the terms containing *a* on one side, and everything else on the other side: $a^2 - 3a - a^2 + ab = -3b - b^2$.

This can be simplified to $-3a + ab = -3b - b^2$.

Take out the common factors *a* on the LHS and -b on the RHS to give: a(b-3) = -b(b+3)

Finish off by

1) a) Express *a* in terms of *b*, given that b(a + 2) = 4.

2) a) Make *x* the subject of the formula $y = \frac{2x^2 - 3}{x^2 - 1}$.



b) Make c the subject of $f = \frac{9}{5}c + 32$.

b) Make *t* the subject of $s = \frac{\sqrt{t+u}}{u}$



already had some practice at it. already had some practice at it.

dividing both sides by
$$(b-3)$$
 to get $a = \frac{-b(b+3)}{b-3}$

Factorising Quadratics

Quadratics have a **Specific Form**

Quadratics are of the form $y = ax^2 + bx + c$, where *a*, *b* and *c* are **numbers**.

A quadratic equation can't have any other powers of x, like x^3 or $x^{\frac{1}{2}}$.

You'll need to be really confident with guadratics for A-level —

they crop up **everywhere**, including in later work on **polynomials** and **graphs** of functions.

Factorising Quadratics is Easier when a = 1

- Normally, you'll factorise your quadratic by splitting it up into 1) two sets of brackets — $x^2 + bx + c = (x + p)(x + q)$.
- 2) As well as factorising a quadratic, you might be asked to **solve** the equation. This just means finding the **values** of *x* that make each **bracket** equal 0 (see example below).
- 3) Factorising quadratics is easier when the coefficient of x^2 is 1.
 - Rearrange your equation into the standard format: $x^2 + bx + c = 0$ 1)
 - 2) Write down the **two brackets** with the x's in: (x + y) = x' + y')(x
 - 3) Now you need to find 2 numbers that **multiply** together to give the value 'c' (the constant) but also **add/subtract** to give the value 'b' (the coefficient of x).
 - Put in the +/- signs and make sure they give the right numbers. 4)
 - 5) **Check** this works by **expanding the brackets** to make sure this gives you the original equation.
 - To solve the equation, set each **bracket equal to 0** and solve for *x*. 6)

EXAMPLE

Solve $x^2 + 7x + 12 = 0$.

This is already in the standard format, with b = 7 and c = 12.

Write out the initial brackets: (x)(x) = 0.

You need to find a pair of numbers that multiply to give c = 12 and add together to give b (= 7): $3 \times 4 = 12$ and 3 + 4 = 7 — perfect.

Write these in the brackets, and then fill in the +/- to make 3 and 4 add to give 7: (x + 3)(x + 4) = 0.

Double check that this works by multiplying out the brackets $(x + 3)(x + 4) = x^2 + 4x + 3x + 12 = x^2 + 7x + 12$ 🖌

= If c is positive, then both signs = 1 f c is negative, = 1 f c is neare the same. It c is negative, then the signs will be opposites.

Set the brackets equal to 0 to solve: (x + 3) = 0 so x = -3 and (x + 4) = 0 so x = -4

What do you call a group of equations? A squadratic...

1) Solve the following quadratic equations:

b) $x^2 + 8x + 7 = 0$ c) $x^2 = 2(7x - 20)$ f) $\frac{x^2}{2} + 2x - 6 = 0$ g) $x - 4 - \frac{12}{x} = 0$ a) $x^2 + 3x + 2 = 0$ d) $x^2 - x = 6$ e) $7x = x^2 + 10$ h) $2x^2 - 2x - 4 = 0$ 2) Factorise $x^2 - 2xz + z^2$.

In an experiment, the temperature $T \circ C$ is modelled by $T = -m^2 + 13m - 30$, where *m* is the time in minutes after the start of the experiment. Find both times at which the temperature is 0 °C.

- a' and b' are the a' and x'
 - and x. 'c' is called the constant term.

multiplying or dividing. E.g. dividing $3x^2 + 6x + 3 = 0$ by 3 gives $x^2 + 2x + 1 = 0$. = by 3 gives $x^2 + 2x + 1 = 0$.

This first step might involve

```
For more on expanding
brackets, see page 10.
```

) = 0

Factorising Quadratics

Factorising a Quadratic is Trickier when a ≠ 1

When $a \neq 1$, the basic method is still the same as before, but it's just a bit more **fiddly**.

- 1) When you write down the two brackets, they'll now be of the form (nx)(mx) where *n* and *m* are two numbers that **multiply** to give *a*.
- 2) Now you're looking for two numbers that **multiply** to give *c*, but which also give you *bx* when you **multiply** them by *nx* and *mx*, and then **add / subtract** them.
- 3) Put those two numbers in the brackets, and then choose their **signs** to make it work.

EXAMPLES

Solve $2x^2 + 7x = -6$.

First rearrange the equation into the standard format: $2x^2 + 7x + 6 = 0$

Now, write out the two brackets. The *x* bits need to multiply together to be $2x^2$ so the brackets will be of the form (2x)(x).

Now, find pairs of numbers that multiply to give c (= 6). Number pairs 1×6 and 2×3 both work.

Try these number pairs out in the brackets until you find a pair that gives 7x. You'll need to try each number in 2 positions because the brackets are different.

 $(2x \quad 1)(x \quad 6)$ multiplies to give 12x and x which add/subtract to give 13x or 11x.

 $(2x \quad 6)(x \quad 1)$ multiplies to give 2x and 6x which add/subtract to give 8x or 4x.

 $(2x \quad 2)(x \quad 3)$ multiplies to give 6x and 2x which add/subtract to give 8x or 4x.

 $(2x \quad 3)(x \quad 2)$ multiplies to give 4x and 3x, which add/subtract to give x or 7x.

This means you can fill in the +/– signs so that 4 and 3 add to give 7: (2x + 3)(x + 2).

Check that this works by multiplying it out: $(2x + 3)(x + 2) = 2x^2 + 4x + 3x + 6 = 2x^2 + 7x + 6$

Set the brackets equal to zero and solve: (2x + 3) = 0, so $x = -\frac{3}{2}$ and (x + 2) = 0, so x = -2

Factorise $3x^2 + 4x - 7$.

The brackets will be (3x)(x).

The only possible number pairing is 1 and 7, so try each number in both positions: $(3x \quad 1)(x \quad 7)$ multiplies to give 21x and x which add/subtract to give 22x or 20x.

 $(3x \quad 7)(x \quad 1)$ multiplies to give 3x and 7x which add/subtract to give 10x or 4x.

Finally work out the signs for each set of brackets: $3x^2 + 4x - 7 = (3x + 7)(x - 1)$.

Don't forget to multiply out the brackets to check your answer is correct.



Oh drat... Double drat.... Triple drat... Quad drat it...

1) Solve the following quadratic equations:

a) $2x^2 + 9x + 9 = 0$ b) $5x^2 + 13x + 6 = 0$

- c) $2x^2 = x + 10$ d) $3x + \frac{21}{x} = 16$
- 2) The equation $h = -4t^2 + 2t + 2$ models the height *h* m a paper plane is from the ground, where *t* is time after the plane is thrown in seconds. At what time *t* does the plane hit the ground?



),

nx, mx, bx... bmx?

The Quadratic Formula

The Quadratic Formula Gives Solutions to Quadratic Equations

The solutions to **any** quadratic equation in the form $ax^2 + bx + c = 0$ are given by this formula:

It means you can solve quadratics that are too hard to factorise.

- 1) To find x, you just need to put the values of a, b and c into the formula.
- 2) The \pm sign means you can end up with two solutions in the final stage you replace it with '+' and then '-'.

EXAMPLE

Solve $8x^2 + 3x - 3 = 0$, giving your answers to 3 decimal places.

a = 8, b = 3 and c = -3. So put these values into the formula:

$$x = \frac{-3 \pm \sqrt{3^2 - 4 \times 8 \times - 3}}{2 \times 8} = \frac{-3 \pm \sqrt{9 + 96}}{16} = \frac{-3 \pm \sqrt{105}}{16}$$

So $x = \frac{-3 + \sqrt{105}}{16}$ and $x = \frac{-3 - \sqrt{105}}{16}$ are the solutions —

you should leave your answers in this form if you're asked for exact solutions.

Put these into your calculator to get: x = 0.453 and x = -0.828



The question mentions decimal places, which suggests you'll need the formula to solve the quadratic.

The **Discriminant** tells you about the **Roots** of a **Quadratic**



 $b^2 - 4ac$

The **discriminant** is the bit of the quadratic formula that's **inside** the **square root**: It's used in A-Level Maths to find **how many roots** a quadratic has. The roots are the **values of** *x* where the **graph** of the quadratic **crosses the** *x***-axis**.

- 1) If the discriminant is **positive** $(b^2 4ac > 0)$, this means the quadratic has **two real roots**. The formula will give **two different real number values** of *x* one from adding and one from subtracting the $\sqrt{b^2 4ac}$ bit.
- 2) If the discriminant is **zero**, the quadratic only has **one root** because adding or subtracting 0 will give the same value of *x*.
- 3) If the discriminant is **negative** $(b^2 4ac < 0)$ this means the quadratic has **no (real) roots** because the square root of a negative number is **not** a real number (see page 6).

EXAMPLE

Find the discriminant of $4x^2 + 3x + 1$. How many real roots does $4x^2 + 3x + 1 = 0$ have?

a = 4, b = 3 and c = 1, so the discriminant is $b^2 - 4ac = 3^2 - 4 \times 4 \times 1 = 9 - 16 = -7$

The discriminant is < 0, so $4x^2 + 3x + 1 = 0$ has **no real roots**.

PRACTICE OUESTIONS

Due to the poor quality of previous jokes, this one's been removed...

1) Solve the following quadratic equations. Give your answers to 3 d.p.

b) 6.

b) $6x^2 + x = 4$

c) $-2x^2 = 3x - 4$

2) Find the discriminant of $5x^2 + 7x + 3$. How many real roots does $5x^2 + 7x + 3 = 0$ have?

a) $4x^2 - 7x + 1 = 0$



Completing the Square

Here's How to **Complete The Square** when **a = 1**

Completing the square means rewriting a **quadratic** as a **squared bracket** plus or minus a **number**. It's used at A-Level for **sketching graphs** (see next page) and **solving the equations** of quadratics (see below) and it's also used with the **equations of circles**. This is the method for when *a* (the coefficient of x^2) is 1:

- 1) Start by getting your quadratic into the **standard format**: $ax^2 + bx + c$.
- 2) Write out the initial squared bracket $(x + \frac{b}{2})^2$ so just divide the coefficient of x by 2.
- 3) Now, **multiply out** the squared bracket and **compare** what you get with the **original** quadratic to see what you need to add or subtract.
- 4) Finally, add or subtract the **adjusting number** to make it **match** the original.

EXAMPLE

Rewrite $x^2 + 12x + 16$ in the form $(x + m)^2 + n$.

This quadratic's already in the standard format, so go straight to step 2 and write out the initial squared bracket: $(x + 6)^2$

Multiply this out and compare with the original: $(x + 6)^2 = x^2 + 12x + 36$

This has a 36 at the end, but you need a 16 at the end, so adjust by 16 - 36 = -20

 $(x + 6)^2 - 20 = x^2 + 12x + 36 - 20 = x^2 + 12x + 16$

So the completed square is $(x + 6)^2 - 20$

To find the adjusting number, you can just take the number term you get from expanding the bracket away from 'c'.

The Completed Square can help you Solve a Quadratic

Once you've **completed the square**, you can **solve** a quadratic. All you have to do is **rearrange** the completed square to make *x* the **subject** by following these three steps:

- 1) Move the **adjusting number** to the **other side** of the equation.
- 2) **Square root** both sides **remember** the ' \pm '...
- 3) Rearrange to get *x* on its **own**.

EXAMPLE

Given that $x^2 - 14x + 36 = (x - 7)^2 - 13$, solve $x^2 - 14x + 36 = 0$.

Set the completed square equal to zero to solve $(x - 7)^2 - 13 = 0$

Now, move the adjusting number to the RHS: $(x - 7)^2 = 13$

Next, square root both sides to get: $x - 7 = \pm \sqrt{13}$

Finally, add seven to each side to get *x* on its own: $x = 7 + \sqrt{13}$ and $x = 7 - \sqrt{13}$

Complete Trafalgar Square — 1 Column + 4 plinths + 5¹⁰ pigeons...

1) Write each of the following in the form $(x + m)^2 + n$:

a) $x^2 + 4x + 1$ b) $x^2 - 12x + 5$

 $x^2 - 12x + 5$ c) $x^2 - 20x - 10$

d) $x^2 + 7x - 3$

States and the second s

E Remember, when you take a square root you get two values —

one positive and one negative.

2) Write $x^2 + 6x - 8$ in the form $(x + m)^2 + n$. Hence solve $x^2 + 6x - 8 = 0$.

You can **Sketch** the **Graph** Using the **Completed Square**

The completed square form of a quadratic gives you information about its graph. When the coefficient of x^2 is **positive**:

- The graph is **u-shaped** with a **minimum point** (see p.37).
- The **adjusting number** tells you the *y*-value of this minimum point. This is because the **smallest value** the quadratic can take occurs when the bit in the brackets is **zero** (the bracket is squared so can never give a negative value).
- If the **adjusting number** is **positive** then the graph's **minimum point** is **above** the *x*-axis. So the graph **never** crosses the *x*-axis it has **no real roots**.

When the coefficient of x^2 is **negative** (there are examples where *a* isn't 1 on the next page):

- The graph is **n-shaped** with a **maximum point** (see p.37).
- The **maximum point** occurs when the bit in brackets is **zero** and the **adjusting number** tells you the *y*-value of this maximum point.
- If the **adjusting number** is **negative** then the graph's **maximum point** is **below** the *x*-axis. So the graph **never** crosses the *x*-axis, meaning it has **no real roots**.

As you saw on the last page, the completed square can also be used to find the solutions of the quadratic. So it can tell you where the graph **crosses the** *x***-axis** (if it crosses the *x*-axis at all).

EXAMPLE

Sketch the graph of $y = x^2 - 14x + 36$.

You know from the example on the previous page that $x^2 - 14x + 36 = (x - 7)^2 - 13$.

The coefficient of x^2 is positive, so it is u-shaped and has a minimum point.

The graph takes its minimum value when $(x - 7)^2 = 0$, so when x = 7. The minimum *y*-value is the adjusting number (-13).

So the minimum point of the graph is at (7, -13).

The graph crosses the *x*-axis at the solutions to the quadratic equation. From the example on the previous page, you know these solutions are $x = 7 + \sqrt{13}$ and $x = 7 - \sqrt{13}$.

To find where the graph crosses the *y*-axis, let x = 0: $0^2 - (14 \times 0) + 36 = 36$, so it crosses at (0, 36).





1)

Complete Times Square — Billboards + billboards + billboards + billboards + billboards + ...

- a) Write $x^2 + 16x + 3$ in the form $(x + a)^2 + b$.
- b) Hence solve $x^2 + 16x + 3 = 0$.
- c) Use this information to sketch the graph of $y = x^2 + 16x + 3$.
- 2) The graph on the right shows the graph of $y = (x + p)^2 + q$, where *p* and *q* are integers.
 - a) Use the sketch to find the values of p and q.
 - b) Find the *y*-intercept of the graph.





Completing the Square

Here's How to **Complete The Square** where **a** ≠ **1**

If *a* isn't 1, you need to take a factor of *a* from the x^2 and x terms before you can complete the square.

This works for negative or This works for negative or fractional values of 'a' as well.

EXAMPLE

Write $3x^2 - 8x = -4$ in the form $p(x + q)^2 + r = 0$.

Start by putting the quadratic into the standard format: $3x^2 - 8x + 4 = 0$.

Take out a factor of 3 from the x^2 and x terms: $3(x^2 - \frac{8}{3}x) + 4 = 0$. Now, write out the initial bracket: $3(x - \frac{8}{6})^2 = 3(x - \frac{4}{3})^2$

Multiply out the bracket and compare to the original: $3(x - \frac{4}{3})^2 = 3x^2 - 8x + \frac{16}{3}$

The adjusting number is
$$4 - \frac{16}{3} = -\frac{4}{3}$$
:
 $3(x - \frac{4}{3})^2 - \frac{4}{3} = 3x^2 - 8x + \frac{16}{3} - \frac{4}{3} = 3x^2 - 8x + 4$

So the completed square is $3(x - \frac{4}{3})^2 - \frac{4}{3}$

Completing the Square is useful for Circle Equations



At A-Level, you'll sometimes need to rearrange the equation of a circle so you can find its **centre** and **radius**. There's lots more about the equations of circles on page 39. With circles you might need to complete the square **twice** — once for x and once for y.

EXAMPLE

Write $x^2 + y^2 - 6x + 14y + 22 = 0$ in the form $(x - m)^2 + (y - n)^2 = k$.

Start by rearranging the equation so the *x*'s and *y*'s are grouped together: $x^2 - 6x + y^2 + 14y + 22 = 0$

Now you need to complete the square for both *x* and *y*. Completing the square for $x^2 - 6x$ gives $(x - 3)^2 - 9$, so the equation so far is: $(x - 3)^2 - 9 + y^2 + 14y + 22 = 0$

Completing the square for $y^2 + 14y$ gives: $(y + 7)^2 - 49$, so the equation is: $(x-3)^2 - 9 + (y+7)^2 - 49 + 22 = 0$

Finally, rearrange into the form $(x - m)^2 + (y - n)^2 = k$: Combine all the number terms together: $(x - 3)^2 + (y + 7)^2 - 36 = 0$ Put the number term on the other side of the equation: $(x - 3)^2 + (y + 7)^2 = 36$



Barney and Pip were meant to be completing the square, but they kept going round in circles.

Complete Red Square — Kremlin + fine this joke's getting old now...

c) $5x^2 - x - 3$

1) Complete the square for each of these quadratics: a) $2x^2 - 3x - 10$ b) $-x^2 - 3x - 1$ 2) Write $x^2 + y^2 - 2x + 6y - 15 = 0$ in the form $(x - m)^2 + (y - n)^2 = k$.

)
$$-\frac{1}{2}x^2 - 3x - 7$$

ď

Algebraic Fractions

Simplify Algebraic Fractions by Cancelling

To simplify an **algebraic fraction**, you need to find **common factors** in the numerator and denominator. Then you can **divide** the numerator and denominator by the common factors to cancel them. Some fractions won't cancel straight away — you'll need to **factorise** first.

Simplify $\frac{36ab^5}{27a^2b^3}$.

The common factors in the numerator and denominator are 9, *a* and *b*³. Cancel down by dividing the numerator and denominator by each one: $\frac{26ab^{5}}{27a^{2}b^{2}} = \frac{4b^{2}}{3a}$

Simplify $\frac{8x^3 - 4x^2}{6x^2 + 5x - 4}$.

Factorising the expressions in the numerator and denominator gives: $\frac{4x^2(2x-1)}{(2x-1)(3x+4)}$

Then cancelling (2x - 1) gives: $\frac{4x^2(2x-1)}{(2x-1)(3x+4)} = \frac{4x^2}{3x+4}$

Multiplying and Dividing Algebraic Fractions

To **multiply** algebraic fractions, multiply the numerators and denominators **separately**. To **divide** algebraic fractions, turn the second fraction **upside down** and then **multiply**.

EXAMPLES

Simplify $\frac{x+3}{2x} \times \frac{x^2}{(x+3)^2}$. Cancel any common factors first: $\frac{x+3}{2x} \times \frac{x^2}{(x+3)^2}$

Then multiply the numerator and denominator separately: $\frac{x}{2(x+3)}$



Simplify $\frac{x^2 - 16}{8} \div \frac{(x - 4)}{2x}$. Start by flipping the second fraction upside down to give: $\frac{x^2 - 16}{8} \times \frac{2x}{(x - 4)}$ Factorise and cancel the common factors (x - 4) and 2: $\frac{(x - 4)(x + 4)}{8} \times \frac{2x}{(x - 4)}$ Then multiply to get the answer: $\frac{x(x + 4)}{4}$



Algebraic Fractions

Adding and Subtracting Algebraic Fractions is a Little Tougher

Before you add or subtract algebraic fractions, take a look at the **denominators**. If they're all the **same**, they have a **common denominator**, so you can just add up the numerators.

But when the denominators are **different**, you'll need to:

- 1) Find a **common denominator** this should be the **lowest common multiple** of all the denominators.
- 2) **Rewrite** each fraction with the common denominator by **multiplying** the top and bottom by the **same thing**.
- Then make into one fraction by adding or subtracting the numerators. 3)

EXAMPLES

Simplify
$$\frac{3x+y}{2} - \frac{x-2y}{7}$$
.

The LCM of 2 and 7 is 14, so rewrite each fraction with a denominator of 14. Multiply the top and bottom of the first fraction by 7, and the top and bottom of the second fraction by 2:

$$\frac{3x+y}{2} - \frac{x-2y}{7} = \frac{7(3x+y)}{14} - \frac{2(x-2y)}{14} = \frac{21x+7y-2x+4y}{14} = \frac{19x+11y}{14}$$

Simplify
$$\frac{1}{x} + \frac{3}{x+1}$$
.

Multiply *x* and (x + 1) together to get the common denominator x(x + 1). Rewrite each fraction with a denominator of x(x + 1) to get:

 $\frac{1}{x} + \frac{3}{x+1} = \frac{1(x+1)}{x(x+1)} + \frac{3x}{x(x+1)} = \frac{x+1+3x}{x(x+1)} = \frac{4x+1}{x(x+1)}$

You can use exactly the same method to add and subtract more than two fractions.

EXAMPLE

Simplify $\frac{1}{4} + \frac{1}{2x} - \frac{1}{4x}$.

The LCM of the three denominators is 4x — so this is the simplest common denominator. Rewriting each fraction over 4x gives: $\frac{x}{4x} + \frac{2}{4x} - \frac{1}{4x} = \frac{x+1}{4x}$



 $\frac{2}{x} + \frac{5a}{x} + \frac{2a^2}{x} = \frac{2+5a+1}{x}$

- These skills are really useful = algebraic fractions pop = up in both years of A-Level.

Inequalities

Solving Inequalities is Just Like Solving Equations...

...whatever you do to one side, you also do to the other side. And follow this important rule:





Quadratic Inequalities are a bit Different

- 1) **Quadratic inequalities** have a **squared term** in them like quadratic equations.
- Quadratic equations can have two solutions (see p.16). Quadratic inequalities are similar — their solutions can be two sets of values, or a range enclosed by two values.

Solve $4x^2 - 14 \le 2$. Solve $3x^2 - 22 > 5$. Add 14 to both sides: $4x^2 - 14 + 14 \le 2 + 14$ Add 22 to both sides: $3x^2 - 22 + 22 > 5 + 22$ $3x^2 > 27$ $4x^2 \le 16$ Divide both sides by 4: $4x^2 \div 4 \le 16 \div 4$ Then divide both sides by 3: $3x^2 \div 3 > 27 \div 3$ $x^2 > 9$ $x^2 \le 4$ You can check your solutions have the right inequality = So $-2 \le x \le 2$ So *x* < −3 or *x* > 3 You can check your solutions have the right inequality Ξ = signs by trying some values. E.g. to check x < -3, try x = -4: $x^2 = 16$ which is > 9, so < is correct. $= try x = -4; x^{2} = 16 \text{ which is } 2, so < is contact. -$

Inequalities

Sketch a Graph to Help You Solve Quadratic Inequalities

When a **quadratic inequality** also has *x*-terms (e.g. $ax^2 + bx + c < d$), it's a bit trickier to solve. The best method is to turn the inequality into an **equation** you can **sketch** — see p.37 for more on sketching quadratics. Then you can solve the inequality by looking for the part of the graph that satisfies it.

EXAMPLE



PRACTICE OUESTIONS 1) Solve a) 4 -2) Use a

I'm no Picasso — but I can sketch a mean quadratic...

1) Solve the following linear inequalities: a) $4 - 8x \le -10x - 6$ b) $-13 \le 2x - 10x - 6$

b) $-13 \le 2x - 3 < 11$ c) -11 < 1 - 3x < 7

- 2) Use a number line to find the values of x which satisfy both 2x 4 < 3x and $5 \ge 7x 2$.
- 3) Solve the following quadratic inequalities: a) $3x^2 - 8 < 4$ b) $2x - x^2 + 8 > 0$ c) $6x \ge 2x^2 + 4$ d) $-5x^2 + 4x + 2 < 6x - x^2$
- 4) Find the values of x which satisfy both $x^2 + 3x + 7 < 3 2x$ and 2(3 2x) > 14.

Graphical Inequalities

Follow These Four Easy Steps to Show Inequalities on a Graph

1) Write each inequality as an **equation**.

Replace the **inequality sign** with an **= sign** and write in the form "y = ..." if you need to.

2) Draw the **graph** of each equation.

Use a **dotted line** if the inequality had a < or > sign and a **solid line** if it had $a \le or \ge sign$.

3) Decide **which side** of each line is the correct one.

Use the origin if possible.

Put the coordinates of a point into each inequality. If the inequality is **true**, the point is on the **correct side** of the line.

4) **Shade** or **label** the correct region.

Check carefully what you're asked to do in the question.

EXAMPLE

Draw and shade the region that satisfies all three inequalities below.

 $3x < 6 + y \qquad -2 + y \le x$

1) Write each inequality as an equation in the form 'y = ':

y = 3x - 6 y = x + 2 y = -x

2) Then draw the graphs. They're all linear equations, so they're just straight lines:



 $y \ge -x$



Graphical Inequalities

You can Show Quadratic Inequalities on a Graph

At A-Level, you'll need to show **quadratic inequalities** on a graph as well as linear ones. To do this, you'll have to accurately sketch a **quadratic graph** using its x- and y-intercepts and turning point. Have a look at p.37 for how — the examples below will just tell you the key points.

EXAMPLES

Draw and label the region that satisfies the inequalities $y > x^2 - 4$ and $3x + 1 \ge y$.

Write the inequalities as equations in terms of *y*:

 $y = x^2 - 4 \qquad \qquad y = 3x + 1$

Then draw the graphs. The quadratic graph is u-shaped. It intersects the *y*-axis at y = -4, and the *x*-axis at x = 2 and x = -2. The turning point is at (0, -4).

Plug the origin (0, 0) into each inequality:

 $y > x^2 - 4 \Rightarrow 0 > -4$ which is true. So the origin is on the correct side of the line.

 $3x + 1 \ge y \Rightarrow 1 \ge 0$ which is true. So the origin is on the correct side of the line.

Label the region above $y = x^2 - 4$ and below y = 3x + 1.



Make sure you get the dotted and solid lines right.

Draw and shade the region satisfying the following inequalities:

$$+ 3x - 2 > y \qquad x^2 < y + 9$$

Write each inequality as an equation in the form 'y = ':

 $-x^2$

$$y = -x^2 + 3x - 2$$
 $y = x^2 - 9$

Then draw the graphs. $y = -x^2 + 3x - 2$ is n-shaped. It intersects the *y*-axis at y = -2 and the *x*-axis at x = 1 and x = 2. The turning point is at (1.5, 0.25).

 $y = x^2 - 9$ is u-shaped. It intersects the *y*-axis at y = -9 and the *x*-axis at x = 3 and x = -3. The turning point is at (0, -9).

Find the region by substituting (0, 0) into each inequality.

 $-x^2 + 3x - 2 > y \Rightarrow -2 > 0$ which is false. So the origin is on the wrong side of the line.

 $x^2 < y + 9 \Rightarrow 0 < 9$ which is true. So the origin is on the correct side of the line.

Shade the region below $y = -x^2 + 3x - 2$ and above $y = x^2 - 9$.





Or cook the inequalities a roast dinner — that'll satisfy 'em...

d)

1) Draw and shade the regions that satisfy the following inequalities:

a)
$$y > x^2$$
 and $8x + 4 < 2y$

c)
$$8 - 2x^2 - y < 0$$
 and $6x > 3y - 18$

b)
$$-x^2 + 4 \ge y$$
 and $y + 10 < 14 - 10$

$$x^2 + 4y > 0$$
 and $x^2 - 6 < y - 2$

2) Draw and shade the region that satisfies the following inequalities: $y < 3 + 5x - 2x^2$ $y \ge x^2 - x - 2$ х

Simultaneous Equations

Use **Elimination** when the Simultaneous Equations are **Linear**

- To solve a pair of simultaneous equations, you need to find the values of 1) the **variables** (e.g. *x* and *y*) which will make **both** equations **true**.
- 2) For example, the simultaneous equations x + 2y = 5 and 3x y = 1 have the solution x = 1and y = 2 because plugging these values into the LHS of the equations gives the RHS.

Here's the **method** for solving simultaneous equations if both equations are **linear**:

- 1) Rearrange the equations into the form ax + by = c.
- 2) Make the **coefficients** of one of the variables the same by **multiplying** one (or both) of the equations by something.
- could make them 4 and -4. 3) Add or subtract the equations to get rid of the terms with the same coefficient, then **solve** to find the value of the **remaining variable**.
- 4) Plug this value back into one of the **original equations** to find the **other variable**.
- 5) Put the values of **both variables** into the **other original equation** to check your answer.

EXAMPLE

Solve the simultaneous equations 3x - 2y = 1 and 2x + 3y = 11.5.

Both equations are already in the form ax + by = c. Label them (1) and (2).

$$3x - 2y = 1$$
 (2) $2x + 3y = 11.5$

Make the coefficients of one of the variables the same. Label the new equations (3) and (4):

$$\begin{array}{cccc} 1 \times 3 & \rightarrow & 9x - 6y = 3 \end{array} \\ \hline 2 \times 2 & \rightarrow & 4x + 6y = 23 \end{array}$$

 $\begin{pmatrix} 1 \end{pmatrix}$

Now eliminate the y's to find the value of x by adding the two equations together:

$$(3) + (4) \qquad 13x = 26 \Rightarrow x = 2$$

Putting this value of x into either equation will allow you to solve for y.

$$a = 2 \text{ in } 2 \text{ gives } 2x + 3y = 11.5$$

$$4 + 3y = 11.5$$

$$3y = 7.5 \Rightarrow y = 2.5$$

Now check your answer by putting both these values into equation (1).

$$x = 2$$
 and $y = 2.5$ in (1) gives $(3 \times 2) - (2 \times 2.5) = 1$

So the solution is x = 2, y = 2.5.

Ling wished she could use elimination to solve all of her problems.

///////////////////////////////////////	11/2
If the signs are the same, then subtrac	t. =
If the signs are opposites, then add.	_
<u> </u>	117

Eliminating the x's — seems a tad cruel...

Solve the following simultaneous equations:

a) y = 4x - 1

5x - 2y = 5

b) 3x + 8y = 2512x - 10y = 16 c) 3x - 2y = 8 d) $\frac{1}{2}x + 2y = 12$ 7x - 5y = 17

y = 4x - 11

Section 4 — More Algebra



S In step 2. vou can ignore the

In step 2, you can ignore the

= signs of the coefficients. E.g. you =

Simultaneous Equations

Use Substitution When One Equation is Non-Linear

If there are any squared terms in the equations, you can't use the elimination method — you have to use the **substitution** method instead. You might have seen this method at GCSE, but you'll use it at **A-Level** with trickier equations, e.g. with more than one squared variable.

- 1) Rearrange the **linear equation** to get a variable **on its own**.
- 2) **Substitute** this into the quadratic equation.
- 3) Rearrange into the **standard format** for a quadratic equation $(ax^2 + bx + c = 0)$ and solve (see p.16-18).
- 4) Plug these values back into the **linear equation** to find the values of the **other variable**.
- 5) Put the values of **both variables** into the **quadratic equation** to check your answer.

EXAMPLE

Label the equ

Solve the simultaneous equations 2y = 5 - x and $x^2 + y^2 = 10$.

Rewrite the linear equation so either *x* or *y* is by itself on one side of the equation.

ations 1 and 2.
1
$$x = 5 - 2y$$

2 $x^2 + y^2 = 10$

Substitute (1) into (2) to form another equation — call this (3).

$$x^{2} + y^{2} = 10$$

(5 - 2y)² + y² = 10 3

Then rearrange (3) to make a quadratic equation in the form $ax^2 + bx + c = 0$ and solve it.

$$25 - 10y - 10y + 4y^{2} + y^{2} = 10$$
 (3)

$$5y^{2} - 20y + 15 = 0$$

$$y^{2} - 4y + 3 = 0$$

$$(y - 3)(y - 1) = 0, \text{ so } y = 3 \text{ or } y = 1$$

You should factorise to solve the quadratic at this step if possible — you can use the quadratic formula if not.

To find the corresponding values of *x*, put each *y*-value back into the linear equation.

Substitute
$$y = 3$$
 into (1) to give: $x = 5 - (2 \times 3) = 5 - 6 = -1$
Substitute $y = 1$ into (1) to give: $x = 5 - (2 \times 1) = 5 - 2 = 3$
Check your answer by putting both pairs of values into equation (2).
 $x = -1$ and $y = 3$ in (2) gives $(-1)^2 + 3^2 = 10$ \checkmark
 $x = 3$ and $y = 1$ in (2) gives $3^2 + 1^2 = 10$ \checkmark
So the solutions are $x = -1$, $y = 3$ and $x = 3$, $y = 1$.

Why did the chickens simultaneously cross the road*...

1) Solve the following simultaneous equations: a) y = 4x + 4 and $y = x^2 + 3x - 8$ c) $x^2 + xy - 10 = 0$ and y + 2x = -7

```
b) y^2 - x^2 = 0 and 3x - y = 3
```

11111111

* [EDIT - don't worry, we've since had words with the 'Editor' responsible for that bad joke...]

Proof

x is just an integer = (a whole number). =

71111111111111111

Remember These Important Facts to Help You Prove Things

You came across proof at GCSE — but at A-Level you'll have to prove some **trickier** things. So you need to make sure you're really comfortable with the **basic skills**. These facts are a good starting point in a lot of proofs:

- 1) You can write any **even number** as 2*x*.
- 2) You can write any **odd number** as 2x + 1.
- 3) You can write **consecutive numbers** as x, x + 1, x + 2, etc.
- 4) You can show something is a **multiple** of a number, x, by showing it can be written ' $x \times$ something'.
- 5) When you **add**, **subtract** or **multiply integers**, you'll always end up with an **integer**.

EXAMPLES

Prove that the difference between any two even numbers is always even.

Write the two even numbers as 2x and 2y, where x and y are integers. Now find the difference: 2x - 2y = 2(x - y). The difference between two integers is an integer, so (x - y) is an integer.

2x - 2y can be written as 2n, where n = x - y. This means that the difference between any two even numbers is a multiple of two, and is therefore even.

Prove that, for any integer *a*, the expression (3a - 1)(a + 2) + 2(a + 3)(a + 2) is always a multiple of 5.

Multiply out the brackets: $3a^2 + 5a - 2 + 2(a^2 + 5a + 6) = 3a^2 + 5a - 2 + 2a^2 + 10a + 12$ Combine like terms: $5a^2 + 15a + 10$ Take out a factor of 5: $5(a^2 + 3a + 2)$. As *a* is an integer, $a^2 + 3a + 2$ is also an integer.

The expression can be written as 5x, where $x = (a^2 + 3a + 2)$, so it is a multiple of 5.

You can Use Facts to Build Up your Argument in a Proof

In some proofs, you'll need to use **facts** and **laws** from maths to show something is true.

EXAMPLE

Prove that the result of dividing a rational number by any other rational number is also rational.

```
You'll learn more about this type of proof at 
A-Level — it's called 'proof by deduction'.
```

Call the two rational numbers *x* and *y*. Rational numbers can be written as fractions with integers on the top and bottom, so write $x = \frac{a}{b}$ and $y = \frac{c}{d}$ where *a*, *b*, *c* and *d* are integers, and *b*, *c* and *d* are not 0 (because you can't divide by 0). $x \div y = \frac{a}{b} \div \frac{c}{d} = \frac{a}{b} \times \frac{d}{c} = \frac{ad}{bc}$. *ad* and *bc* are the products of two integers, so they are also integers. *bc* is not 0, as neither *b* nor *c* is zero. **The result of dividing two rational numbers can be written as a fraction with**

integers on the top and bottom and a non-zero denominator, so it is also rational.



Using proof by deduction, Ama worked out that her sister was the jumper-stealing culprit.
Proof

You can **Disprove** by Finding a **Counter Example**

Finding a **counter example** is one of the easiest ways to prove a statement is **wrong**. If you can just show **one case** of where the statement doesn't work, then you've disproved it. Sorted.

EXAMPLES

Find a counter example to disprove the following statement: "When x is prime, 2x + 1 is prime."

Try different primes in the expression.

Substituting x = 3 into 2x + 1 gives $(2 \times 3) + 1 = 7$ 7 is a prime number, so try again...

Substituting x = 7 into 2x + 1 gives $(2 \times 7) + 1 = 15$ 15 is not prime, so the statement is **not true**. It can sometimes take a while to find a counter-example, but just keep your cool...

By finding a counter example, disprove the following statement: "Given that *a* and *b* are integers, if a > b, then $a^2 + 1 > b^2 + 1$."

At first glance, this might look correct — but take a closer look at the statement. *a* and *b* are integers, which means they can be negative numbers too...

So if a = 1 and b = -2, then 1 > -2, so a > b. But $a^2 + 1 = 2$ and $b^2 + 1 = 5$, so $a^2 + 1 < b^2 + 1$.

For a = 1 and b = -2, the second part of the statement isn't satisfied.

So this statement is **not true**.

Ian says, "If x and y are two different irrational numbers, then xy must also be irrational." Find an example to show that Ian is wrong.

The square roots of any non-square number are irrational, so try $x = \sqrt{8}$ and $y = \sqrt{2}$:

 $xy = \sqrt{8} \times \sqrt{2} = \sqrt{8 \times 2} = \sqrt{16} = 4$

4 is not an irrational number, so lan is wrong.

You saw on p.6 that an irrational number is one that can't be written exactly as a fraction.

PRACTICE QUESTIONS

I looked for the proof in my pudding — but I just got really sticky...

- 1) Prove that $(n + 1)^2 (n 1)^2$ is even for any integer *n*.
- 2) Prove that the product of any two odd numbers is always odd.
- 3) Prove that the result of subtracting a rational number from any other rational number is also rational.
- 4) Phyllis believes the following statement is true: " $2^n + 1$ always gives a prime number, when *n* is an integer." Prove, by finding a counter example, that Phyllis is wrong.
- 5) Find a counter example to disprove the following statement: "For any pair of integers x and y, $\frac{x^2}{y^2} < \frac{x}{y}$."

Functions Take an Input and Give an Output

1) A **function** is a type of **mapping** — a set of **instructions** on how to get from one number to another. There are two main ways to write a function:

f(x) = 8x + 1
$f: x \to 8x + 1$

These both mean that 'f' takes a value for x_i , **multiplies** it by 8, **adds** 1, and **outputs** the result.

3) The set of values you put into a function is called the **domain**. The set of values output from a function is called the **range**. A function maps **each value** in the **domain** to **one** value in the **range**.

You can **evaluate** a function for different values by substituting each value into the function: 4)

Functions

EXAMPLES

2)

Find f(4) for the function $f(x) = (x + 6)^2$.

Just put x = 4 into the function and work out the answer: $f(4) = (4 + 6)^2 = 10^2 = 100$

 $f: x \rightarrow 8x - 7$. Find f(-3). Substitute x = -3 into the function: $f(-3) = (8 \times -3) - 7 = -31$

You can **Put Functions Together** to Make **Composite Functions**

1) When you have **two different functions**, say f(x) and g(x), you can combine them to make a **single new function**. This is called a **composite function**.

- 2) The composite function of f(x) and g(x) is written fg(x) — which means 'do g first, then apply f to your answer'. ff(x) or $f^2(x)$ means 'do f **twice**'.
- 3) You **always** do the function **closest to** *x* first.

EXAMPLES

 $f(x) = x^2 + 20$ and g(x) = 2x + 3, find: a) fg(x) $fg(x) = f(g(x)) = f(2x + 3) = (2x + 3)^2 + 20$

$$gf(x) = g(f(x))$$

= g(x² + 20) = 2(x² + 20) + 3 = **2x² + 43**

ATTATION 111111 Usually, you'll find that $fg(x) \neq gf(x)$. 7111111111111111

 $f(x) = x^2 + 1$, g(x) = 4 - 2x and h(x) = 3 + x. Find fgh(3).

 $(or 4x^2 + 12x + 29)$

Start with the function closest to x and work outwards. So do h(3) first: 3 + 3 = 6. Then apply g to this result: $g(6) = 4 - (2 \times 6) = -8$

Then apply f to this result: $f(-8) = (-8)^2 + 1 = 65$, so fgh(3) = 65.

 $x^2 = 4 \Rightarrow x = -2 \text{ or } x = 2$

 $f(x) = 2x^2 - 1$. Find the value(s) of x when f(x) = 7.

 $2x^2 - 1 = 7$

 $2x^2 = 8$

Set the function equal to 7 and solve:

b) gf(x)

= There's more about manning There's more about mappings, domains and ranges in the second year of A-Level Maths. = - second year of / Loron /

Composite functions were covered at GCSE, but there's = some trickier stuff to learn about them in Year 2 of A-Level — so make sure you know the basics here.

You might not have seen this notation at GCSE, so don't panic if you don't recognise it.



Functions

An Inverse Function Reverses the Effect of the Original Function

- 1) The **inverse** of a function **reverses** what the function does. The inverse of a function f(x) is written with the notation $f^{-1}(x)$.
- 2) You'll see inverse functions in more detail in Year 2 of A-Level.
- 3) You can **find the inverse** of a function using these steps:
 - 1) Write out the function 'f(x) = ...' but **replace** f(x) with x, and substitute y into the expression instead of x. This gives you x = f(y).
 - 2) **Rearrange** to make *y* the **subject**.
 - 3) Then **replace** y with $f^{-1}(x)$.

EXAMPLE

If f(x) = 4x + 12, find $f^{-1}(x)$.

Replace f(x) with x and x with y: x = 4y + 12

Then rearrange to make *y* the subject: $4y = x - 12 \Rightarrow y = \frac{x - 12}{4}$

Finally, replace y with $f^{-1}(x)$ to get: $f^{-1}(x) = \frac{x-12}{4}$

If you **combine** a function f(x) with its **inverse** $f^{-1}(x)$, you get x. $f^{-1}f(x) = ff^{-1}(x) = x$ This is because f(x) and $f^{-1}(x)$ just **reverse** each other, so the **output** is the **original input**. You can use this to check whether you've worked out an inverse function correctly.

EXAMPLE

a) If f(x) = 8 - 12x, find $f^{-1}(2)$.

Find $f^{-1}(x)$ — replace f(x) with x, and x with y: x = 8 - 12y

Rearrange for *y*: $y = \frac{8-x}{12}$

Replace *y* with $f^{-1}(x)$ to get $f^{-1}(x) = \frac{8-x}{12}$ Substitute *x* = 2 into the function: $\frac{8-2}{12} = \frac{1}{2}$ b) Show that $ff^{-1}(x) = x$ for this function.

You know that $f^{-1}(x) = \frac{8-x}{12}$, so just put this into f(x) and simplify.

$$ff^{-1}(x) = f\left(\frac{8-x}{12}\right) = 8 - 12\left(\frac{8-x}{12}\right)$$
$$= 8 - (8-x) = x$$
$$= \frac{3}{5} ff^{-1}(x) = x, \text{ so the inverse}$$

z is correct. Hoorah. z



So if f(2) = 5, then $f^{-1}(5) = 2$.

"Inverse <u>fun</u>ctions?" cried Mark, "but those sound like the opposite of fun!"

Straight Lines

The Equation of a Straight Line is y = mx + c

You need to know this stuff like the back of your hand for A-Level it comes up all over the place, from **logarithms** to **differentiation**. If you know the **gradient** and the *y*-intercept of a straight line, you can give its **equation**.

> The equation of any straight line can be written in the form y = mx + cwhere *m* is the gradient of the line and *c* is the *y*-intercept of the line.

c is the *y*-intercept because it's the point on the graph where x = 0 — so $y = m \times 0 + c = c$. Similarly, to find the *x*-intercept you set *y* equal to 0 and solve for *x*.

EXAMPLE

A straight line has a gradient of 4 and a *y*-intercept of 6. Find the point where this line crosses the *x*-axis.

First find the equation of the line. The gradient m = 4 and y-intercept c = 6, so y = 4x + 6. Now set y equal to zero and solve for x: $0 = 4x + 6 \Rightarrow 4x = -6 \Rightarrow x = -1.5$. So the line crosses the x-axis at the point (-1.5, 0).

You can also read off the gradient and *y*-intercept from the equation of a line. At **A-Level**, you'll have to deal with straight-line equations written in **different forms**, e.g. ax + by + c = 0, rather than y = mx + c. Just do a bit of rearranging to get the equation into the form you want.

EXAMPLE

Find the gradient of the line 3x + 4y + 4 = 0.

Rearrange into y = mx + c form: $4y = -3x - 4 \Rightarrow y = -\frac{3}{4}x - 1$. So the gradient is $-\frac{3}{4}$.

You can Find the Gradient using Two Points on a Line

You can take any two points on a line and use their coordinates to find the gradient of that line.

The gradient between two points (x_1, y_1) and (x_2, y_2) is given by change in $x = \frac{y_2 - y_1}{x_2 - x_1}$

EXAMPLE

Find the gradient of the straight line on the right.

Pick two accurate points on the line and label them (x_1, y_1) and (x_2, y_2) . Here, you can read off the points $(x_1, y_1) = (-2, 3)$ and $(x_2, y_2) = (4, -2)$.



Plug these into the formula to find the gradient: $\frac{-2-3}{4-(-2)} = -\frac{5}{6}$



1) Find the gradient, *x*-intercept and *y*-intercept of the line 2x + y = -2.

- 2) A line has gradient 3 and *y*-intercept 5. Find the equation of the line.
- 3) Find the gradient of the line that goes through the points (3, 4) and (-2, 1).

35

Straight Lines

Find the Equation of a Line using Two Points on the line

If you know two points that a line goes through, you can work out the equation of the line.

EXAMPLE

Find the equation of the line that passes through points (1, 7) and (3, 16). Give your answer in the form ax + by + c = 0 where a, b and c are integers.

Label the points: $(x_1, y_1) = (1, 7)$ and $(x_2, y_2) = (3, 16)$

Find the gradient with $\frac{y_2 - y_1}{x_2 - x_1}$: $\frac{16 - 7}{3 - 1} = \frac{9}{2}$, so $y = \frac{9}{2}x + c$.

Substitute one of the points into the equation you've found and solve for *c*:

$$y = \frac{9}{2}x + c \Rightarrow 7 = \frac{9}{2} + c \Rightarrow c = 7 - \frac{9}{2} = \frac{5}{2}$$
. This gives $y = \frac{9}{2}x + \frac{5}{2}$

Multiply the whole equation by 2 to get integer coefficients: 2y = 9x + 5. Then rearrange into the correct form: 9x - 2y + 5 = 0

Find the Distance between Two Points with Pythagoras



Pythagoras' theorem tells you that, for any **right**angled triangle, $a^2 + b^2 = c^2$. So length $c = \sqrt{a^2 + b^2}$. At A-Level you'll use this to calculate the magnitude of vectors...

You can use this to find the **distance** between **two points** (x_1, y_1) and (x_2, y_2) :

- 1) Use the two points to make a **right-angled triangle**, with the **hypotenuse** as the line between the two points.
- The difference between the *x*-values will give the length of one side. The difference between the *y*-values will give the length of the other side.

Distance between (x_1, y_1) and $(x_2, y_2) = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$



EXAMPLE

Point A is (4, 17) and point B is (10, 2). What length is line AB?

Label the points: $(x_1, y_1) = (4, 17)$ and $(x_2, y_2) = (10, 2)$, so the length of AB is given by $\sqrt{(10-4)^2 + (2-17)^2} = \sqrt{6^2 + (-15)^2} = \sqrt{261} = 16.16$ (2 d.p.)



You can do a surprising amount with two points...

- 1) Find the equation of the line that passes through points (1, 2) and (5, 9).
- 2) A line is drawn between (-3, 4) and (7, -2). Find:
 a) The equation of this line in the form *ax* + *by* + *c* = 0 where *a*, *b* and *c* are integers.
 b) The length of the line. Give your answer to 2 d.p.
- 3) The distance Moe walks is modelled by a straight line graph. At time t = 2 hours, he has walked a distance of d = 10 kilometres. At time t = 3.5 hours, he has walked a distance of d = 17.5 kilometres.
 - a) Give the equation of the line in the form d = mt + c.
 - b) How long does it take Moe to travel 12.5 km?

Parallel and Perpendicular Lines

At A-Level, you'll need these skills in parts of the differentiation topic, as well as in coordinate geometry.

Parallel Lines have Equal Gradient

Parallel lines have the same gradient, which means they never cross each other.

EXAMPLE

Line A has equation y = 3x - 2. Line B passes through point (4, 3) and is parallel to line A. Find the equation of line B.

Parallel lines have the same gradient, so gradient of line B = gradient of line A = 3.

This gives you the first bit of the equation for line B: y = 3x + c.

You know the point (4, 3) is on the line, so plug this into the equation and solve for c: $3 = 3 \times 4 + c \Longrightarrow 3 = 12 + c \Longrightarrow c = 3 - 12 = -9.$

So the equation of line B is y = 3x - 9.

Perpendicular Lines Cross at Right Angles

The gradients of perpendicular lines multiply together to give -1. So:

The gradient of the perpendicular line = $-1 \div$ the gradient of the other line

EXAMPLE

Find the equation of the line perpendicular to 3y - 2x + 4 = 0 that passes through (2, 2). Give your answer in the form ax + by + c = 0, where *a*, *b* and *c* are integers.

First, rearrange the equation into y = mx + c form: 3y = 2x - 4 so $y = \frac{2}{3}x - \frac{4}{3}$.

The perpendicular line will have gradient $-1 \div \frac{2}{3} = -1 \times \frac{3}{2} = -\frac{3}{2}$.

So the equation so far is: $y = -\frac{3}{2}x + c$.

Plug in the point (2, 2) and solve for *c*: $2 = -\frac{3}{2} \times 2 + c$

 $2 = -3 + c \Longrightarrow c = 2 + 3 = 5$

So the equation of the perpendicular line is y = -x + 5.

Finally, rearrange into the form asked for in the question.

Multiply by 2 to get rid of the fractions: 2y = -3x + 10.

Then put everything on one side of the equation: 3x + 2y - 10 = 0.



Parallel lines meet about as often as the bad guy wins in Bond films...

- Line F has equation 5y + 3x 7 = 0. Line G is parallel to line F and goes through the origin. What is the equation of line G?
- 2) Lines P and Q are perpendicular and meet at the point (-1, -2). The gradient of line P is 4. Find the equations of P and Q. Give your answers in the form ax + by + c = 0.
- The points P (4, 3) and Q (9, 5) lie on line l_1 . The line l_2 is perpendicular to l_1 and 3) passes through Q. Find the equation of line l_2 in the form ax + by + c = 0.

36



Chiara lived in a parallel

universe where she couldn't be told off for crossing the line.

NIIIIIIIIIIIIIIIIIIII You might see perpendicular lines called



divide by a fraction, you flip it _____ over and multiply (see p.7).

(3, 0)

37

All Quadratic Graphs have a Symmetrical Bucket Shape

You'll need to be really confident with **sketching quadratics** at A-Level — there's a whole topic on it. You'll also use these skills in the **inequalities** and **kinematics** topics. VIIIIIIIIIIIIIIIII To remember, think = To remain for negative...

All **quadratic** graphs have the **same shape** — a **symmetrical curve**.



Negative quadratics (ones where the coefficient of x^2 is negative) are **n-shaped** like this.

Work Out Three Things to Sketch A Quadratic

If you're asked to **sketch** a quadratic, you **don't** need to use graph paper and carefully plot every point. You just need to work out and label the **most important** bits of the graph to make sure your drawing is **roughly correct**. As well as the shape of the graph, you need to work out:

- 1) Where the graph **crosses** the *y*-axis (the *y*-intercept).
- 2) Where the graph **crosses** the *x*-axis (the *x*-intercepts). These are called the **roots** of the quadratic.
- 3) You might be asked to find the **turning point** (or 'vertex') of the graph — this could be the **minimum** or **maximum** point depending on whether it's **u-shaped** or **n-shaped**.



(0, 6)

0

(2.5, -0.25)

EXAMPLE

Sketch the graph of $y = x^2 - 5x + 6$, including any points of intersection with the axes and the vertex of the graph.

The coefficient of x^2 is positive, so the graph is **u-shaped**.

To find the *y*-intercept, let x = 0: $y = 0^2 - 5 \times 0 + 6 = 6$. So it crosses the *y*-axis at y = 6.

The graph intersects the *x*-axis when y = 0, so solve $x^2 - 5x + 6 = 0$ by factorising: $x^{2} - 5x + 6 = (x - 2)(x - 3) = 0$, so the *x*-intercepts are x = 2 and x = 3.

Quadratic graphs are symmetrical, so the *x*-coordinate of the turning point of the graph is halfway between 2 and 3. So it's $(2 + 3) \div 2 = 2.5$.

Plug this number back into the quadratic to find the y-coordinate of the turning point: $2.5^2 - 5 \times 2.5 + 6 = -0.25$. So the turning point is (2.5, –0.25).

Now you can sketch the graph. Make sure to label each of the points.





Symmetrical bucket hats — the must-have maths fashion trend...

1) Sketch these quadratics, labelling the points of intersection with the axes and the vertex of each graph. a) $y = x^2 - 3x$ b) $y = x^2 + x - 2$ c) $y = -x^2 + 6x - 9$ d) $v = 3x^2 + 2x - 8$



Harder Graphs

A **Cubic** Contains an **x**³ Term

Cubics are of the form $ax^3 + bx^2 + cx + d$ where a, b, c and d are all numbers ($a \neq 0$). Cubic graphs have a characteristic 'wiggly' shape.

If x^3 has a **positive** coefficient, then the graph goes from the **bottom** left to the top right of the axes.



If x^3 has a **negative** coefficient, then the graph goes from the top left to the bottom right.



At A-Level you'll learn how to factorise and sketch simple cubics. Here's an example of how to sketch one.

EXAMPLE

Given that $x^3 + 3x^2 - x - 3 = (x + 1)(x - 1)(x + 3)$, sketch the graph of $y = x^3 + 3x^2 - x - 3$.

The method for sketching a cubic is similar to the method for sketching a quadratic.

The x^3 term has a positive coefficient, so the graph will go from the bottom left to the top right.

Now you just need to find where the graph crosses the axes. It crosses the *y*-axis when x = 0, so at y = -3. It crosses the *x*-axis when y = 0: $x^3 + 3x^2 - x - 3 = (x + 1)(x - 1)(x + 3) = 0$, so it crosses the x-axis at x = -1, 1 and -3.

Now you can use these values to sketch the graph.



on exponentials at A-Level. Graphs of the form $y = k^x$ or $y = k^{-x}$ (for any positive number *k*) are called **exponential** graphs. An exponential graph is a **curve** that is always **above** the *x***-axis**. All exponential graphs go through the point (0, 1) — because anything to the power 0 is 1.





What goes up must come down — unless it's an exponential graph...

Sketch the graphs of the following cubics: a) y = (x + 1)(x - 1)(x + 2)b) y = (x - 1)(x - 2)(x - 3)

c) $y = -x^3 - 5x^2 + 6x$

Section 5 — Graphs



,11111111111111

You'll cover lots



Harder Graphs

Reciprocal Graphs have Equation **y** = **k**/**x** or **xy** = **k**

- 1) Reciprocal graphs have **two curves** in **diagonally opposite quadrants**. They're **symmetrical** about the lines y = x and y = -x.
- 2) The pair of quadrants the curves are in depends on whether *k* is **positive** or **negative**.



When *k* is **negative**, the graph looks like this.



- 3) **Reciprocal graphs** are undefined at x = 0 (the *y*-axis) and y = 0 (the *x*-axis).
- 4) So reciprocal graphs **never touch** the axes but they do get **infinitely close**. This means the axes are **asymptotes** of the graph you'll be expected to know this term at A-Level.



Circles have Equations $(x - a)^2 + (y - b)^2 = r^2$

The general equation for a **circle** with **centre** (*a*, *b*) and **radius** *r* is $(x - a)^2 + (y - b)^2 = r^2$. Circle equations of the form $x^2 + y^2 = r^2$ were covered at GCSE. These circles have centre (0, 0), so are the **special case** where a = b = 0.

EXAMPLE

2)

Find the equation of the tangent to $(x - 5)^2 + (y - 5)^2 = 50$ at the point (-2, 4).

From the general equation of a circle, you know the centre is (5, 5). Start by finding the gradient of the radius — the line from the centre of the circle to the point (-2, 4).

Gradient of radius =
$$\frac{\text{change in } y}{\text{change in } x} = \frac{5-4}{5-(-2)} = \frac{1}{7}$$

A tangent to a circle meets a radius at 90°, so they are perpendicular (this is one of the circle theorems from GCSE).

This means the gradient of the tangent is $-1 \div \frac{1}{7} = -7$.

So the equation of the tangent so far is y = -7x + c.

Substitute in the point (-2, 4) and solve for *c*: $4 = -7 \times -2 + c \Rightarrow 4 = 14 + c \Rightarrow c = -10.$

So the equation of the tangent is y = -7x - 10.



Stilling See page 36 for more on perpendicular lines.

Reciprocal graphs — a good way to get one over on your x...

1) Give the centre and radius of the following circles: a) $(x - 1)^2 + (y - 3)^2 = 16$ b) $x^2 + (y + 2)^2 = 50$

c)
$$(x + \frac{1}{2})^2 + (y - \frac{1}{2})^2 = 2$$

Find the equation of the tangent to the circle $(x - 1)^2 + (y - 2)^2 = 8$ passing through point (3, 4).

Graph Transformations

f(x) + a is a Translation Along the y-axis

A transformation of the form f(x) + a is a **translation parallel** to the *y*-axis. The graph of f(*x*) moves *a* units in the *y*-direction (i.e. vertically), so all of the *y*-values of f(*x*) will have *a* added to them. Translating a graph doesn't change its shape, it just **moves** it.

EXAMPLE

The graph of y = f(x) has a minimum point at (2, -4). Write down the coordinates of the minimum point of f(x) + 5.

This is a translation along the *y*-axis — so it only affects the *y*-values. This means you add 5 onto the *y*-value of the minimum point. So the minimum of f(x) + 5 is (2, -4 + 5) = (2, 1).

At A-Level, you might have to describe a translation using a **column vector**.

This type of translation can be described by the vector $\begin{pmatrix} 0 \\ a \end{pmatrix}$.

f(x + a) is a Translation Along the x-axis

A transformation of the form f(x + a) moves the graph of f(x)**left** or **right** — i.e. **parallel** to the *x*-axis.

Be **careful** with these — when *a* is **positive**, the graph moves to the **left**. When it is **negative**, the graph moves to the **right**.

EXAMPLE

The diagram shows the graph of y = f(x). Sketch the graph of y = f(x + 3).

a = 3, so the graph moves 3 units in the negative x-direction. This means the graph moves to the left by three units.

Even if you're given a graph that you don't recognise, just remember how the translations work and you'll be fine. Temember how the translations work and you'll be mile.

This type of translation can be described by the column vector $\begin{pmatrix} -a \\ 0 \end{pmatrix}$.





I can't find that graph — it must've been lost in translation...

- 1) The diagram on the right shows the graph of y = f(x):
 - a) Sketch the following translations:
 - i) f(x) + 3, labelling the coordinates of the minimum and where the curve meets the y-axis.
 - ii) f(x 2), labelling the coordinates of the minimum and where the curve meets the x-axis.
 - b) State the column vectors that describe the above translations.



 $(-1, \frac{-16}{3})$



Make sure you're really confident with all these types of transformation — you'll learn a few more at A-Level.

Make sure you're round confident with all these types of transformation — you'll faw more at A-Level.

Paula's home transformation was going to take a lot of hard graph-t.

y = f(x)



Graph Transformations



EXAMPLES

a) $f(x) = x^2 + 2x - 8$. Find the *y*-intercept of y = -f(x).

The *y*-intercept of f(x) is when x = 0. $f(0) = (0)^2 + 2(0) - 8 = -8$, so is at (0, -8).

The *y*-intercept of y = -f(x) is the *y*-intercept of y = f(x) multiplied by -1. So y = -f(x) intersects the *y*-axis at (0, 8).

b) Given that f(x) = (x - 2)(x + 4), find the roots of f(–*x*).

The roots of f(x) are x = 2 and x = -4.

f(-x) is a reflection in the *y*-axis, so all the *x*-values are multiplied by -1. This means the roots of f(-x) are x = -2 and x = 4.

The diagram shows the graph of y = g(x). Sketch the graph of y = -g(x).

The graph of y = -g(x) is a reflection of y = g(x) in the x-axis.

The *x*-values stay the same, but the *y*-values are multiplied by -1. This means the *y*-intercept is 1 instead of -1.





Now is a good time to stop and reflect on how far we've come...

- The diagram on the right shows the graph of y = f(x). 1) Sketch the graphs of y = -f(x) and y = f(-x), labelling the *x*- and *y*-intercepts of each.
- 2) The quadratic graph y = g(x) and its reflection y = g(-x) are identical. For each of the following, state whether it could be g(x). a) $x^2 -$ $-x^2$

4 b)
$$x^2 + 3x - 1$$
 c)

3) y = h(x) is the graph of a quadratic with the turning point (4, -7). State the coordinates of the turning point of the following graphs:

a) y = -h(x)b) y = h(-x)

Trigonometry — Sin, Cos, Tan

opposite

10 m

Use **SOHCAHTOA** to Remember how to use **Sin, Cos** and **Tan**

Remember SOH CAH TOA for working out side lengths and angles in right-angled triangles:

At A-Level, you'll come across a lot more trigonometry, = so you need to have a good grip on the basics.

hypotenuse

 $\sin x = \frac{\mathbf{o}\text{pposite}}{\mathbf{h}\text{ypotenuse}}$ $\cos x = \frac{\mathbf{a}\text{djacent}}{\mathbf{h}\text{ypotenuse}}$ $\tan x = \frac{\mathbf{o}\text{pposite}}{\mathbf{a}\text{djacent}}$

- The **hypotenuse** is the longest side of the triangle.
- The **opposite** is the side opposite the angle you're working with (*x*).
- The **adjacent** is the side next to the angle you're working with (*x*).



EXAMPLE

Calculate the height of this isosceles triangle to 3 s.f.

Start by splitting the triangle down the middle to get two right-angled triangles. Then you can use trigonometry on one of them.



You have the adjacent side and want to find the opposite side, so use tan:

 $\tan 55^\circ = \frac{x}{5} \Rightarrow x = 5 \times \tan 55^\circ = 7.1407... = 7.14 \text{ m (3 s.f.)}$

To calculate a missing angle, you need to use one of the **inverse trig functions** — **sin**⁻¹, **cos**⁻¹ or **tan**⁻¹.

EXAMPLE

Find angle *x* to 3 s.f.

You know the opposite and the hypotenuse, so you need to use sin:



$$\sin x = \frac{5}{7} \Rightarrow x = \sin^{-1}\left(\frac{5}{7}\right) = 45.5846... = 45.6^{\circ}$$
 (3 s.f.)



Trigonometry — Sin, Cos, Tan

Learn these Trig Values

There are **two triangles** that you can use to help you work out important trig values:





```
You can use Pythagoras'
Theorem to check that the
side lengths are right (p.35).
```

Using SOH CAH TOA on these triangles gives you these values:





Sarah found trigonometry difficult, so she decided to work on her tan.

You can Calculate Trig Values using the Unit Circle

The unit circle is a circle with radius 1 centred on the origin.

At A-Level, you'll see the unit circle used to calculate trig values.

- 1) You can take a **point** on the unit circle and make a **right-angled triangle**.
- 2) You know the **hypotenuse** of this triangle will be **1**, as it is the **radius** of the unit circle.



- 3) This means you can find the **coordinates** of any point (*x*, *y*) on the unit circle by using trig the coordinates are given by (**cos** *a*, **sin** *a*).
- 4) The angle *a* needs to be measured **anticlockwise** from the **positive** *x***-axis**.

EXAMPLE

Find the coordinates of point P on the unit circle. Give your answer to 2 d.p.

The point is on the unit circle, so you know the *x*-coordinate is $\cos 130^\circ = -0.64278...$ and the *y*-coordinate is $\sin 130^\circ = 0.76604...$) So the coordinates of P to 2 d.p. are (**-0.64, 0.77**)





Trigonometry — Graphs

The Graph of y = sin x Repeats Every 360°

You should have seen **all three** of the trig graphs at GCSE. You'll need to be really comfortable with them at A-Level, because they're used for **solving trig equations**. This is the graph of $y = \sin x$:



It'd be a sin not to learn every last bit of this page... 1) By sketching a graph, find all the solutions to the following equations in the interval $0^\circ \le x \le 360^\circ$. Where necessary, give your answers to 1 decimal place. a) $\sin x = \frac{1}{\sqrt{2}}$ b) $\sin x = -0.7$ c) $\sin x = -0.2$

Trigonometry — Graphs

The Graph of y = cos x is the Same Shape as sin x

The graph of **cos** x is the **same shape** as the graph of **sin** x, but shifted left by 90°. So the graph of $y = \cos x$ looks like this:



Because it's the same shape as the sin graph, it shares some of the same properties:

- 1) The *y*-values for the graph are all **between –1 and 1**.
- 2) The basic shape of the graph **repeats every 360°**, so $\cos x = \cos (x + 360°) = \cos (x 360°)$.

But there are some differences between the two:

- 1) $\cos x$ is **symmetrical** about the *y*-axis you can **reflect** it in the *y*-axis and it'll look the same. So $\cos(-x) = \cos(x)$.
- 2) Because $\cos 0^\circ = 1$, the graph crosses the *y*-axis at y = 1.
- 3) It crosses the *x*-axis at \pm 90°, \pm 270° etc.

EXAMPLE



Solve $\cos x = -0.8$ for $-180^\circ \le x \le 180^\circ$. Give your answers to 1 d.p.

Solving $\cos x = -0.8$ with a calculator will give you $x = \cos^{-1}(-0.8) = 143.1301...^{\circ}$

This gives one answer. Now sketch the graph over this interval to find the second solution.

As every 360° interval of the graph is symmetrical, the second solution will be the same distance to the left of the *y*-axis as $^{-1}$ the first is to the right. The first solution is 143.1301...° to the right of the *y*-axis, so the second solution is 143.1301...° to the left of the *y*-axis. So x = -143.1301...°

So the solutions to $\cos x = -0.8$ for $-180^{\circ} \le x \le 180^{\circ}$ are $x = 143.1^{\circ}$ and $x = -143.1^{\circ}$ (1 d.p.)



Cosimodo — The humpback trig graph of Notre Dame...

1) By sketching a graph, find all the solutions to the following equations in the interval $-360^\circ \le x \le 360^\circ$. Where necessary, give your answers to 1 decimal place.

a) $\cos x = \frac{1}{\sqrt{2}}$

- b) $\cos x = 0.1$
- c) $\cos x = -0.4$

Just like with sin x, if the first solution you find is outside the interval, add or subtract multiples of 360° to find a solution inside the interval.

Trigonometry — Graphs

The Graph of y = tan x Repeats Every 180°

The graph of $y = \tan x$ is completely different from the other two. It looks like this:



- 1) Unlike the graphs of sin and cos, the graph of $y = \tan x$ can give **any value** of y. The graph takes **every** y value between $-\infty$ and $+\infty$ in each **180° interval**.
- 2) The graph **repeats every 180°** and goes through the origin (because $\tan 0^\circ = 0$).
- 3) The graph is **not defined** at $\pm 90^{\circ}$ and at $\pm 270^{\circ}$:
 - These points are marked on the graph by dotted lines called asymptotes.
 - As the graph approaches each asymptote from the **left**, it shoots **up to infinity**.
 - As it approaches from the **right**, it shoots **down to minus infinity**.
 - The graph **never** actually **touches** the asymptotes, although it does get **infinitely close**.

EXAMPLE

Solve $\tan x = -5$ for $0^{\circ} \le x \le 360^{\circ}$. Give your answer to 2 d.p.

Use your calculator to get a solution to the equation: $\tan^{-1}(-5) = -78.69...$ This is outside the interval, so add 180° to find the first solution: -78.69...° + 180° = 101.309...°Sketch the graph of $y = \tan x$ to find

Sketch the graph of $y = \tan x$ to find the solutions within the given range.

You can see from the graph that the next solution will be 180° + 101.309...° = 281.309...°



So the solutions are *x* = 101.31° and *x* = 281.31° (2 d.p)



Fake tan x — it's not quite orange, but it gets infinitely close...

) By sketching a graph, find all the solutions to the following equations in the interval $-360^\circ \le x \le 360$. Where necessary, give your answers to 1 decimal place. a) tan x = -1 b) tan x = 7 c) tan x = -6

The Sine and Cosine Rules

The Sine and Cosine Rules work for Any Triangle

- To work out side lengths and angles in **any triangle** (not just right-angled ones), you'll need to use the **sine** and **cosine rules**. These came up at GCSE, but you need to be slick at using them for A-Level.
- 2) Be **careful** how you **label** the triangle when using the sine and cosine rules — the side **opposite** an angle should have the **same letter**. E.g. side *b* is opposite angle *B*.

The Sine Rule $\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$





Which rule you use depends on which sides and angles you know in the triangle.

Learn when to use the Sine Rule

If you have **two angles** and a **side**, you use the **sine rule** to find the lengths of the other sides. Because you have two angles, you can find the **third** by **subtracting** them from **180**°.

EXAMPLE





When Sam saw the cloud, he knew it must be a sine.

If you're given two sides and an angle that's not between them, you need to use the sine rule.

EXAMPLE

Find angle *BAC* in the triangle shown.

The known angle is not enclosed by the two known sides, so use the sine rule. Find angle *C* first:

$$\frac{b}{\sin B} = \frac{c}{\sin C} \Rightarrow \frac{11}{\sin 41^{\circ}} = \frac{7}{\sin C}$$

Rearrange to get sin
$$C = \frac{7 \times \sin 41^{\circ}}{11} = 0.417... \Rightarrow C = \sin^{-1}(0.417...) = 24.676...^{\circ}$$

This means that angle $BAC = 180^{\circ} - 41^{\circ} - 24.676...^{\circ} = 114.3^{\circ}$ (1 d.p.)



11 cm

The Sine and Cosine Rules

Learn when to use the Cosine Rule

If you're given **two sides** and the **angle between them**, you can use the **cosine rule** to find the length of the third side.

EXAMPLE

Two ships set sail from the same port. Ship X travels 42 miles due north. Ship Y travels 36 miles on a bearing of 035°. How far apart are the two ships?

Draw a diagram and fill in the known values.

Label the sides and angles -z is the side you're trying to find.

From the diagram, you can see that the two known lengths are either side of the known angle, so you need to use the cosine rule.

 $z^2 = 36^2 + 42^2 - 2 \times 36 \times 42 \times \cos 35^\circ = 582.884...$

So $z = \sqrt{582.884...} = 24.1$ miles (1 d.p.)



Area = $\frac{1}{2}ab\sin C$

If you know **all three sides** of the triangle but no angles, then you need the **cosine rule**.

EXAMPLE



There's also a Formula for the Area of a Triangle

The **area** of a triangle is given by this formula: To use the **formula**, you need **one angle** and the **two sides enclosing the angle**. If you don't have all of these, you'll need to use the sine or cosine rule to find them.

EXAMPLE

Find the area of this triangle to 1 d.p.

You're given an angle and two sides enclosing it, so you can plug the numbers straight into the formula.





Vectors

Vectors have a Direction and a Magnitude

You will have met **vectors** at GCSE, but you're going to see a whole lot more of them at A-Level, especially in **mechanics**. They're often used for modelling things like forces.

- 1) Scalars are quantities without a direction e.g. a speed of 5 m/s. They're just numbers.
- Vectors represent a movement of a certain magnitude (size) in a direction

 e.g. if two objects have a velocity of 5 m/s and -5 m/s, this means
 they are travelling at the same speed but in opposite directions.

There are **Different Ways** to **Represent Vectors**

- 1) Vectors are **drawn** as **arrows**, like this: 3 m/s
 - The **direction** of the vector is shown by the **arrowhead**.
 - The **size** of the vector is shown by the line's **length**.
- 2) Vectors are written as a lowercase **bold** letter (**a**) or a lower case **underlined** letter (<u>a</u>).
- 3) If the **start** and **end point** of a vector are known, it might also be written like this: \overrightarrow{AB} .
- 4) Vectors can also be written as column vectors A
 - e.g. $\begin{pmatrix} 4 \\ 1 \end{pmatrix}$ is a vector that goes **four units right** and **one unit up**.

Adding Vectors Describes Movements Between Points

- 1) Adding two vectors $\mathbf{a} + \mathbf{b}$ means 'go along \mathbf{a} then go along \mathbf{b} '.
- 2) To add two vectors you can draw their arrows **nose to tail**.
- 3) The **single** vector that goes from the **start** of the first vector in the sum to the **end** of **the last** is called the **resultant vector**.
- 4) **Subtracting** one vector from another is a little more complicated — because $-\mathbf{b}$ is just the vector **b** reversed, you can think of $\mathbf{a} - \mathbf{b}$ as $\mathbf{a} + (-\mathbf{b})$. So $\mathbf{a} - \mathbf{b}$ means 'go along **a** then backwards along **b**'.

EXAMPLE

Find \overrightarrow{AD} in terms of s, t and u.

The first thing to do is relabel the vectors so they're nose to tail from A to D like this.

Now you can see that to get from A to D, you go along **-s**, **t** and then **-u**:

 $\overrightarrow{\mathsf{AD}} = -\mathbf{s} + \mathbf{t} + (-\mathbf{u}) = \mathbf{t} - \mathbf{s} - \mathbf{u}$





Woe to the scalars — and to the vector go the spoils... U 1) $\overrightarrow{AB} = 2\mathbf{s} + \mathbf{t}$ and $\overrightarrow{BC} = 2\mathbf{t} - \frac{1}{2}\mathbf{s}$.

D

Show that $2\overrightarrow{AC} = 3(\mathbf{s} + 2\mathbf{t})$.

2) Using the diagram on the right, find \overrightarrow{TW} in terms of the vectors **a** and **b**.

BĂ

R



The second se

helps with vector questions.

At GCSE you may

have also seen vectors

written like this — a.



ĀB

۱A/

Vectors

Multiplying a Vector by a Scalar Changes its Size

- 1) Multiplying a vector by a **positive scalar** changes the vector's **size**, but **not** its direction.
- Multiplying a vector by a negative scalar changes the vector's size and its direction gets switched.
- 3) To multiply a **column vector** by a **scalar**, multiply the top and bottom numbers **separately** by the scalar, e.g. $2 \times \begin{pmatrix} 4 \\ 1 \end{pmatrix} = \begin{pmatrix} 2 \times 4 \\ 2 \times 1 \end{pmatrix} = \begin{pmatrix} 8 \\ 2 \end{pmatrix}$.
- 4) To show two vectors are parallel, you need to show they are scalar multiples of each other.

EXAMPLE

 \overrightarrow{XQ} = b and \overrightarrow{PQ} = a. Point P lies halfway along \overrightarrow{XY} and point Q lies halfway along \overrightarrow{XZ} . Show that \overrightarrow{YZ} is parallel to \overrightarrow{PQ} . As Q is the midpoint of \overrightarrow{XZ} , \overrightarrow{XZ} = $2\overrightarrow{XQ}$ = 2b.

As Q is the midpoint of XZ, $XZ = 2XQ = 2\mathbf{b}$. $\overrightarrow{XP} = \mathbf{b} - \mathbf{a}$. P is the midpoint of \overrightarrow{XY} , so $\overrightarrow{XY} = 2\overrightarrow{XP} = 2(\mathbf{b} - \mathbf{a})$.



You can use **Vectors** to Show **Three Points** are on a **Line**

- 1) If three or more points all lie on a single straight line, they are collinear.
- 2) To show three points A, B and C are collinear, you need to show that \overrightarrow{AB} and \overrightarrow{BC} are parallel.
- 3) So you need to show that the vectors are **scalar multiples** of one another.

EXAMPLE

Show that X, Y and Z are collinear.

First, find \overrightarrow{XY} and \overrightarrow{YZ} : $\overrightarrow{XY} = -(\mathbf{d} - \mathbf{c}) + \mathbf{d} = \mathbf{c}$ and $\overrightarrow{YZ} = -\mathbf{d} + \mathbf{d} + 2\mathbf{c} = 2\mathbf{c}$. $\overrightarrow{YZ} = 2\overrightarrow{XY}$, which means \overrightarrow{XY} and \overrightarrow{YZ} are parallel, so X, Y and Z are collinear.

A Vector's Magnitude is its Length





b) $\begin{pmatrix} -1 \\ -1 \end{pmatrix}$

At **A-Level** you'll have to calculate the **magnitude of a vector**.

Magnitude of vector $\begin{pmatrix} a \\ b \end{pmatrix} = \sqrt{a^2 + b^2}$

$$a^2 + b^2$$
 b



Getting from point A to point C has never been so exciting...

1) Show that the vectors $\mathbf{v} = 4\mathbf{a} + 6\mathbf{b}$ and $\mathbf{u} = 6\mathbf{a} + 9\mathbf{b}$ are parallel.

2) Using the diagram on the right, show that W, X and Y are collinear.

3) Calculate the magnitude of the following vectors: a) $\begin{pmatrix} 3 \\ 2 \end{pmatrix}$

 $N \xrightarrow{2\mathbf{u} - 4\mathbf{v}}_{X} \xrightarrow{\mathbf{v}}_{\mathbf{v}}$

Sampling

A Sample Tells You About a Population

In A-Level Maths, you'll cover sampling in more detail and learn some new sampling methods. Here's a **recap** on what sampling is, with a couple of new terms thrown in.

- 1) The **population** is the **whole group** you want to investigate.
- 2) If you collect information from every single member of the population, it's called a census. A census gives really accurate results, but it's usually too difficult or time-consuming to survey everyone.
- 3) That's where **sampling** comes in you can choose a **selection** from the population to **represent** the whole group. This smaller group is called a **sample**.
- First you need to identify all the people or things you can **sample from** these are 4) the **sampling units**, and a **full list** of the sampling units is called the **sampling frame**. For reliable results, your sampling frame should include as much of the population as possible.

You have to **Choose** your Sample **Carefully**

- 1) It's really **important** that a sample is **similar** to the population. This means that the sample is **representative** of the population, which allows you to use it to draw conclusions about the whole population.
- 2) A **biased** sample is one that **doesn't** fairly represent the population. It can be hard to avoid bias altogether, but there are a few **rules** you can follow to **limit it** as much as possible.
- Choose from the **correct population** and don't exclude anyone.
- Choose your sample at random.
 - Make sure your sample is **big enough**.
- You need to be able to **spot sampling bias** think about **when**, 3) where and how the sample is taken, and if it's big enough.

Choose a **Random** Sample using **Simple Random Sampling**

One way to get a **random** sample is to use the **simple random sampling** method. This gives every member of the population an **equal chance** of being in the sample, so it's **unbiased**. To **select** a simple random sample, start by giving **every** member of the population a **unique** number. Then generate a list of random numbers and match them to the population members.

EXAMPLE

A retailer has 700 store card holders. Describe how simple random sampling could be used to select a sample of 30 of the card holders to survey.

Start by drawing up a list of all the card holders, and give each one a 3-digit number between 001 and 700.

Next, generate a list of 3-digit random numbers, using a calculator or random number table. You should reject any numbers over 700, or any that are repeated.

meeting as their sample, explain why this sample may be biased.

Choose sample members randomly — it's common census...

A council wants to survey the residents of a town about local issues. They hold a meeting one lunchtime where residents can give their views. If the council uses the residents who attend the

Stop when you have a list of 30 unique numbers, and select the card holders with the matching numbers — this is your sample.







by sampling all of the tarts.









1)

Learn the **Difference** between **Quantitative** and **Qualitative** Data

1) A **variable** is a quantity that can take **different** values. There are two broad **categories**:

Qualitative variables take non-numerical values — e.g. colours of cars. **Quantitative** variables take **numerical values** — e.g. temperature.

There are two different types of **quantitative data** — **discrete** and **co** 2)

Discrete variables take certain values within a particular range — e. Continuous variables can take any value within a particular range –

Get familiar with the basics Get familiar with the basics = - vou'll need them for the

A-Level statistics topics.

You need to Know about **Two Types** of **Frequency**

- 1) **Frequency tables** show how many times a data value occurs.
- 2) Grouped frequency tables group data into classes. These classes tell you the **range** that the value is contained in. The upper class boundaries and lower class boundaries can be defined with inequalities to cover all possible values.
- You can work out the **class width** with this formula: 3)

Class width = upper class boundary – lower class boundary

You can also calculate 4) the mid-point of a class.

upper class boundary + lower class boundary Mid-point =

EXAMPLE

The masses of 21 rocks (to the nearest 10 g) are recorded in the table. Add columns to the table to show the upper and lower class boundaries, class widths and the class mid-points.

The lightest rock that would weigh 150 g to the nearest 10 g is 145 g. So the lower class boundary of the 150-250 class is 145 g.

The upper class boundary of the 150-250 class is 255 g. This is the same as the lower class boundary of the 260-320 class, so there are no gaps between classes.

Find the class boundaries for the other classes, then use the formulas to calculate each class width and mid-point:

Mass (g)	Frequency
150-250	7
260-320	9
330-400	5

The data has been rounded, so the table doesn't show the actual class boundaries.

These classes can be written using

These classes can be written using

1 inequalities. E.g. the 150-250 class

- would be written $145 \leq m < 255$. would be written $145 \le m < 255$.

Mass (g)	Frequency	Lower class b'dary (g)	Upper class b'dary (g)	Class width (g)	Mid-point (g)
150-250	7	145	255	110	200
260-320	9	255	325	70	290
330-400	5	325	405	80	365



What kind of quantitative data do spies use? Discreet variables...

The table on the right shows the length (to the nearest 10 mm) of 50 pieces of string. Add four columns to show the lower class boundaries, upper class boundaries, the class widths and the class mid-points.

Length (mm)	Frequency
80-110	15
120-200	16
210-320	19

ontinuous	A-Level sta	atistics topics.
.g. shoe size – e.g. length		
Table		
	Mass (m, g)	Frequency
	$150 \le m < 180$	2

 $180 \le m < 250$

 $250 \le m < 300$

lower class boundary

12 upper class boundary

6

Histograms Show Frequency Density

You'll have seen histograms at GCSE — but they rear their ugly heads again at A-Level.

1) **Histograms** are like bar charts, but the **height** of each bar tells you the **frequency density** instead of the frequency. You can calculate frequency density with this **formula**:

Frequency Density = Frequency ÷ Class Width

Histograms are for continuous data.

- 2) The **width** of the bars tells you the **class widths**.
- 3) The **area** of each of the bars tells you the **frequency**. This is because you can **rearrange** the **formula** for frequency density to give: **Frequency = Frequency Density × Class Width**.

EXAMPLE

a) The table and histogram show information about the masses of sharks in an aquarium. Use the histogram to find the missing entry in the table.

First, you'll need to add an extra column to the table to calculate frequency density:

Now, use those frequency densities to work

 $1050 < m \le 1130$

out the scale of the vertical axis of the graph:

This means you can read off the remaining bar -

it has a frequency density of 0.3, and the table shows a

class width of 80, so the missing frequency is $0.3 \times 80 = 24$.

24

Mass (m, kg)	Frequency	Frequency Density
$750 < m \leq 850$	10	$10 \div 100 = 0.1$
$850 < m \leq 900$	12	$12 \div 50 = 0.24$
$900 < m \leq 1050$	51	$51 \div 150 = 0.34$
$1050 < m \le 1130$		

Mass (<i>m</i> , kg)	Frequency
$750 < m \le 850$	10
$850 < m \le 900$	12
$900 < m \le 1050$	51
1050 < m < 1130	

Frequency density



b) Estimate the number of sharks that weigh between 800 kg and 950 kg.

Use the fact that frequency = frequency density × class width to estimate the frequency between 800 kg and 950 kg. The class widths are the bits of the classes you're interested in: Between 800 and 850 kg, there are an estimated $0.1 \times (850 - 800) = 0.1 \times 50 = 5$ sharks. Between 850 and 900 kg, there are 12 (you can read this from the table). Between 900 and 950 kg, there are an estimated $0.34 \times (950 - 900) = 0.34 \times 50 = 17$ sharks.

0.3

This makes the estimate 5 + 12 + 17 = 34 sharks that weigh between 800 kg and 950 kg.



Those who don't learn from histograms are doomed to repeat stats...

1) This table shows information about the heights of a group of people.

- a) Draw a histogram to represent the information.
- b) Estimate the number of people between 155 cm and 200 cm tall.

_			
	Height (h, cm)	Frequency	
	$150 < h \leq 170$	40	
	$170 < h \leq 190$	32	
	$190 < h \le 210$	26	
	$210 < h \leq 240$	9	

Averages

The Mode and Median are Averages

You'll need to know all about the **different types** of **average**.

The **mode** is the most **commonly occurring** data value.

The **median** is the **middle value** of the data, when the data is sorted in **order of size**.

Before you calculate the median, you need to put your **data** in **ascending order**. Then you need to work out the **middle position**.

- 1) If you have an **odd** number of data values, then the **middle** point will be a **data value**.
- If you have an **even** number of data values, the **middle** will be **between** two data values. 2) So your **median** will be **halfway between** those values.

EXAMPLE

Find the median of this set of data: 8, 12, 4, 6, 13, 7, 8, 4.

This data is modes are 4 and 8. First, put the data in ascending order: 4, 4, 6, 7, 8, 8, 12, 13. There are 8 data values and $8 \div 2 = 4$. So the median will be halfway between the fourth and fifth data values in the list. These values are 7 and 8, so the median is 7.5.

The Mean is the Sum of the Data Divided by the Total Fi

The mean is another type of average. At A-Level you'll use some new notation f Don't worry though — you still calculate it like you did at GCSE.

1) The notation for the mean is \overline{x} (said 'x-bar'). The **formula** is given as:

- 2) Each *x* is a **data value**, *f* is the **frequency** of each *x* and *n* is the **total number** of data values.
- The \sum (called sigma) just means you **add things together**. So $\sum x$ is just the **sum** of all the **data values** x. 3) $\sum f$ is the sum of the frequency of the data values — it's the same as saying 'the total number' of data values.
- So to find the mean of 1, 7, 3 and 5, add them all together to get 16 and then 4) divide by 4 (because there are four values). This gives a mean of 4.

You can calculate the **combined mean** of **two data** sets using this **formula**: All you need to know is the **mean** of each data set $\overline{x_1}$ and $\overline{x_2}$, and the **size** of each data set n_1 and n_2 .

EXAMPLE

The mean of a set of 5 numbers is 20. The mean of another set of 7 numbers is 13. Find the combined mean of both sets of numbers.

Plug these numbers into the combined mean formula:

$$\overline{x} = \frac{5 \times 20 + 7 \times 13}{5 + 7} = \frac{191}{12} =$$
15.92 (2 d.p.)



Statistical Nursery Rhyme Idea #12 — Eeny Meany Median Mode...

- 1) Find the mode, median and mean of this data set: 1, 7, 8, 5, 4, 6, 5.
- 2) The mean of a set of 5 numbers is 12. The mean of another set of 6 numbers is 15. Find the combined mean of both sets of numbers.

For it.

$$\sum x = \sum fx$$

$$\overline{x} = \frac{\sum x}{n}$$
 or $\frac{\sum fx}{\sum f}$



At A-Level, you may see these averages called 'measures of location'.

If a set of data has two modes then it is called bimodal

then it is called bimodal. 71111111111111111111111111

Estimate Averages for Grouped Data

- 1) The **modal class** is the class with the highest **frequency density**.
- 2) You can also find the **class containing the median** at A-Level you'll learn how to estimate the median.
- To estimate the mean, you'll need to add extra **columns** one for the 3) **mid-point** of each class and another for **frequency** × **mid-point**.
- Then the class frequency density will just class with the highest frequency. Because you don't know **exact** values when you work with **grouped data**, these averages are only **estimations**. 4)

EXAMPLES

This table shows information about the widths of cardboard boxes. State the modal class.

As a	II the cl	ass wie	dths a	re the	same,	the	modal	class	will
be t	he class	with t	he hig	hest fi	requen	cy: 4	40 < w	≤ 60	cm.

a) This table shows information about the length (to the nearest 10 mm) of 60 tree branches. Write down the class containing the median.

There are 60 data values, so the median is halfway between the 30th and 31st values. Count through the 'Frequency' column — both these values lie in the third class, so the class containing the median is 110-160 mm.

b) Calculate an estimate for the mean length.

You'll need to add two extra columns to the table — one for each class's mid-point (x)and one for the value of the frequency multiplied by the class's mid-point (fx). You also need to add a total row.

Each class's mid-point is halfway between the upper and lower class boundaries. Be careful with the first one though the upper class boundary is 55 and

the lower class boundary is 0, so the mid-point is $(0 + 55) \div 2 = 27.5$.

So the total frequency $(\sum f)$ is 60 and the total of the frequencies multiplied by the mid-points $(\sum fx)$ is 9012.5.

Put these into the formula for the mean: $\overline{x} = 9012.5 \div 60 = 150.21$ (2 d.p.).



Learn this and become a lean, mean, average estimating machine...

- 1) This table shows information about the masses of pieces of fruit picked in an orchard (to the nearest 10 g). a) Find the class containing the median.
 - b) Calculate an estimate for the mean weight.

	0
Mass (g)	Frequency
0-40	6
50-90	11
100-140	8
150-190	5

Width (w, cm)	Frequency
$20 < w \leq 40$	10
$40 < w \le 60$	11
$60 < w \le 80$	9

Length (mm)	Frequency
0-50	5
60-110	7
110-160	22
160-210	16
210-260	10

Length (mm)	Frequency (f)	x	fx
0-50	5	27.5	137.5
60-110	7	85	595
110-160	22	135	2970
160-210	16	185	2960
210-260	10	235	2350
Total	60	_	9012.5

21		1111	11111	111	/11//	
= '	it the	class	widths	are	all the	52000
-					an the	same, -

Cumulative Frequency is the Running Total of the Frequency

- 1) You can draw a graph of the cumulative frequency of a data set, which lets you estimate the **median** and the **upper** and **lower quartiles** for the data.
- The upper quartile is the value 75% of the way through an ordered set of data. The lower quartile is the value 25% of the way through an ordered set of data.
- 3) The interquartile range is the difference between the upper and lower quartiles.

EXAMPLE

a) The table below shows information about the masses of 100 puppies. Draw a cumulative frequency graph for the data.

The first thing you need to do is add a third column to the table for cumulative frequency.

To draw the graph, plot cumulative frequency up the vertical axis, and mass along the horizontal axis.

Start by plotting a cumulative frequency of zero at the lowest value in the first class — no puppies weigh less than 50 g.

Plot the rest of the points using the highest value of each class and the cumulative frequency.



Mass (m, g)	Frequency	Cumulative Frequency
$50 \le m < 150$	5	5
$150 \le m < 210$	7	5 + 7 = 12
$210 \le m < 270$	16	12 + 16 = 28
$270 \le m < 330$	25	28 + 25 = 53
$330 \le m < 410$	31	53 + 31 = 84
$410 \le m < 460$	11	84 + 11 = 95
$460 \le m < 500$	5	95 + 5 = 100

b) Use your graph to estimate the median and interquartile range of the weights.

To find the median, go halfway up the side to 50 and read off the value from the curve: so an estimate of the **median** is **325 g**.

To find the interquartile range, you need to find the upper and lower quartiles first. The upper quartile is 75% of the way through the data — so go across from 75 and read off the value from the curve: you get 380 g. The lower quartile is 25% of the way through the data — so go across from 25 and read off the value from the curve: you get 260 g. So an estimate of the **interquartile range** is 380 g – 260 g = **120** g.

terturtle range — about a flipper and a half				
	Length (<i>l</i> , mm)	Frequency		
The table on the right shows information about	$300 \le l < 350$	6		
some lengths of copper wire.a) Draw a cumulative frequency graph for the data.b) Use your graph to estimate the median and the interquartile range of the lengths.	$350 \le l < 410$	10		
	$410 \leq l < 470$	24		
	$470 \le l < 520$	14		
	$520 \le l < 600$	6		

Ir

Probability

Remember these Important Probability Facts

You'll meet trickier probability in A-Level Maths, and build on your knowledge in new topics like **probability distributions**. So make sure you're **happy with the basics** here.

- 1) Finding the **probability** of something means working out **how likely** it is to happen. Your answer should always be between **0** (**impossible**) and **1** (**certain**).
- For any probability experiment, the different things that can happen are called outcomes. E.g. if you flip a coin, the outcomes are 'heads' and 'tails'. The set of all possible outcomes is called the sample space.
- 3) The thing you want to find the probability of is often called an 'event'. An event matches one or more of the possible outcomes, e.g. if you roll a dice, one event is 'roll a number less than 3' — which matches the outcomes '1' and '2'. You can write 'the probability of an event' as 'P(event)' for short.
- 4) There's a handy formula you can use to work out the probability of an event when all the possible outcomes are equally likely:



5) Another useful fact is that probability always **adds up to 1**. So for a set of events where **only one** of them can happen at a time, the probabilities of **all** the possible events **add up to 1**.

You can **Find** Probabilities from **Two-Way Tables**

You might be asked to work out probabilities using **information** in a **two-way frequency table**. The numbers in the table tell you how many **outcomes** match different **events**, and you can find the **total number** of outcomes in the **bottom right hand corner**.

EXAMPLE

a) A bag contains 30 balls — coloured red, purple, or yellow, and labelled A or B. The numbers in each category are shown in the table. Find the probability that a ball selected at random is purple.

	Red	Purple	Yellow	Total
Α	7	3	4	14
В	3	5	8	16
Total	10	8	12	30

There are 8 purple balls in total, so there are 8 outcomes where the event 'ball is purple' happens. The total number of possible outcomes is the total number of balls, which is 30.

So using the probability formula above: P(ball is purple) = $\frac{8}{30} = \frac{4}{15}$

In the question, 'at random' tells you the outcomes are all equally likely, so you can use the formula.

b) Find the probability that a ball selected at random is red or labelled A.

You need the number of red balls, plus the number of purple and yellow balls labelled A

- that's 10 + 3 + 4 = 17. So P(ball is red or labelled A) = $\frac{17}{30}$



Chances are you almost certainly didn't hate this page — right?

1) I roll a fair, six-sided dice. Find the probability that I roll: a) 1 or 2 b) higher than 2.

2) Using the two-way table in the example above, find the probability that a randomly selected ball is yellow and labelled A.



The probability of Hector being a good boy is 1.

You can have a cup of tea venn you've done these questions...

- 1) Out of 40 customers at a book shop, 22 bought fiction books, 7 bought travel books, and 2 bought both fiction and travel books. Draw a Venn diagram to represent the 40 customers.
- Using the Venn diagram in the second example above, find the probability 2) that a randomly selected student likes biology or chemistry or both.

58

A Venn Diagram Shows How Things are Grouped

Venn diagrams show how a **collection of objects** is **split up** into different **groups**.



Probability

Finding Probabilities from Venn Diagrams

You've used Venn diagrams to find probabilities at GCSE, and you'll use them again at A-Level. They're really useful for solving probability problems — the labels show the **number of outcomes** (or the **probabilities**) matching the **different events** that can happen.

EXAMPLE

This Venn diagram shows the number of children at a primary school who go to baking club (B) and singing club (S). Find the probability that a randomly selected child from the school goes to baking club.

The event 'child goes to baking club' is represented by circle B. So there are 15 + 10 = 25 outcomes matching this event. And there are 15 + 10 + 20 + 15 = 60 children in total.

So using the formula from p.57 you get: P(child goes to baking club) = $\frac{25}{60} = \frac{5}{12}$

Here's a trickier example with three circles — and this time the diagram is labelled with the proportion in each area, rather than the number.

EXAMPLE

A group of students were asked which of biology (B), chemistry (C) and physics (P) they like. The diagram shows the proportion who like each subject. If one student is selected at random, find the probability that they like physics, but not chemistry.

'Likes physics, but not chemistry' is shown by the area that's in P, but not in C.

The probability of the student being in this area is equal to the proportion in the area.

So add up the numbers to get: P(likes physics, but not chemistry) = 0.05 + 0.05 = 0.1

В С 0.25 0.25 0.15 0 0.05 0.05



Laws of Probability

The Addition Law Tells You P(A or B)

Say you have **two events**, A and B. You can find the probability of **event A or event B or both** happening using the **addition law**.

If the events A and B **can't happen** at the same time, they're called **mutually exclusive** events. For mutually exclusive events A and B, **P(A and B) = 0**, so the addition law becomes:

You can **show** whether or not two events are **mutually exclusive**. All you have to do is to show whether or not P(A and B) **equals zero**.

EXAMPLE

P(A) = 0.16, P(B) = 0.5 and P(A or B) = 0.62. Show whether or not events A and B are mutually exclusive.

Rearrange the addition law to find P(A and B): P(A or B) = P(A) + P(B) - P(A and B) $\Rightarrow P(A \text{ and } B) = P(A) + P(B) - P(A \text{ or } B) = 0.16 + 0.5 - 0.62 = 0.04$

 $P(A \text{ and } B) \neq 0$, so events A and B are **not mutually exclusive**.

The Product Law Tells You P(A and B)

If one of events A and B happening **doesn't affect** the **probability** of the other happening, **events A and B** are **independent**. This means you can use the **product law** for **independent** events:

This means you can use the **product law** for **independent** events:

EXAMPLE

Sid buys one piece of fruit each day. The probability he buys an apple, P(A), is always 0.4, independent of any other day. Find the probability that he buys an apple on exactly one of the next two days.

This can happen in one of two ways: either Sid buys an apple on the first day and not on the second day, or he doesn't buy an apple on the first day, but does on the second day.

P(A) = 0.4, so P(not A) = 1 - 0.4 = 0.6, since probabilities add up to 1.

The days are independent, so you can multiply the probabilities using the product law to find P(A and 'not A') and P('not A' and A): P(A and 'not A') = $0.4 \times 0.6 = 0.24$ and P('not A' and A) = $0.6 \times 0.4 = 0.24$.

These two options can't both happen — either one happens or the other happens, so you add the probabilities using the addition rule for mutually exclusive events: P(one apple in two days) = P('A and not A' or 'not A and A') = P(A and 'not A') + P('not A' and A) = 0.24 + 0.24 = 0.48.

Like with mutually exclusive events, you can **show** whether or not two events are **independent**. You just need to show whether or not $P(A) \times P(B) = P(A \text{ and } B)$ — see Question 2 below.

PRACTICE OUESTIONS 1) 2)

Probability law #3 — learn probability laws 1 and 2, or else...

1) On any given day, the probabilities that Martha and Sahil play tennis are 0.3 and 0.4 respectively. Find the probability that, on a given day, Martha or Sahil plays tennis.

) If P(A) = 0.6, P(B) = 0.5 and P(A and B) = 0.3, show that events A and B are independent.

You'll see how to write it using set notation in Year 2 of A-Level Maths.

 P(A or B) = P(A) + P(B) - P(A and B)

This law should be familiar from GCSE.

You need a different form of the law for dependent events — it's

- E covered in Year 2 of A-Level.
- $P(A \text{ and } B) = P(A) \times P(B)$

P(A or B) = P(A) + P(B)

Tree Diagrams

Tree Diagrams Show Probabilities for a Sequence of Events

To **draw** a tree diagram representing two 'trials':

- 1) Draw a set of branches for the first trial, with a branch for each result.
- 2) At the **end** of each branch, draw a set of branches for the **second trial**.
- 3) **Label each branch** with the **probability** of that result probabilities on each **set of branches** should **add up to 1**.
- 4) To find the **probability** of a sequence of events, **multiply along the branches**. The probabilities of **all the sequences** added together should **always be 1**.

EXAMPLE

60

A fair, six-sided dice is rolled three times. Use a tree diagram to find the probability that exactly two even numbers are rolled.

There are three trials and you can define two results for each trial — 'even number' (E), and 'odd number' (O), where each result has a probability of $\frac{1}{2}$.

There are three sequences that give two even numbers — (E, E, O), (E, O, E) and (O, E, E). Multiplying along the branches, the probability of each sequence $=\frac{1}{2} \times \frac{1}{2} \times \frac{1}{2} = \frac{1}{8}$.

You want to find the probability of (E, E, O) or (E, O, E) or (O, E, E), so add the probabilities:

P(exactly two even numbers) = P(E, E, O) + P(E, O, E) + P(O, E, E) = $\frac{1}{8} + \frac{1}{8} + \frac{1}{8} = \frac{3}{8}$

Tree Diagrams can show Dependent Events

Events are **dependent** if one event happening **affects the probability** of the other happening. When you're drawing a tree diagram for dependent events, you need to **change** the **probabilities** on the **second** set of branches, **depending** on the result in the **first** set.

EXAMPLE

A cake tin contains 6 chocolate muffins and 4 blueberry muffins. Two muffins are taken at random without replacement. Using a tree diagram, find the probability that at least one chocolate muffin is taken.

There are two trials, each with results 'chocolate' (C) and 'blueberry' (B). The probabilities for the second trial change because the first muffin isn't replaced. There are now 9 muffins left in total, and the number of each flavour depends on the first selection. P(at least one chocolate) = 1 - P(no chocolate) = 1 - P(two blueberry) = 1 - $\left(\frac{4}{10} \times \frac{3}{9}\right) = 1 - \frac{2}{15} = \frac{13}{15}$



Sustainable stats — for every page I write, I draw a tree diagram...

1) Claire eats either pizza or curry every Friday. The probability that she has pizza is 0.25 if she had pizza the previous Friday, and 0.6 if she had curry the previous Friday. Given that Claire has curry this Friday, find the probability that she has pizza on both of the next two Fridays.

A 'trial' is just something like rolling a dice, or selecting an object at random. For three trials, just repeat step 2.

13 a) x = 2 or x = 1

Page 1 — Diagnostic Test

- 1 4, -10, 205 and 0 are integers.
- 2 5.9, $\frac{1}{5}$, -6, $\sqrt{4}$, 13.978 and 2.1 are rational. π and $\sqrt{7}$ are irrational.

3 a)
$$\frac{2}{15}$$

b) $\frac{1}{4}$
c) $\frac{11}{12}$
d) $\frac{51}{35}$
4 a) x^9
b) $2y^2$
c) 1
d) $32n^{10}$
5 $\frac{1}{25}$
6 a) $\frac{9}{16}$
b) 4
c) 4
d) $\frac{1}{6}$
7 a) $x^2 - 2x - 24$
b) $x^2 + 10x + 25$
c) $2x^2 + 5x - 3$
d) $x^3 + 2x^2 - 19x - 20$
8 a) $5(x + 4)$
b) $3a(1 + 4b)$
c) $(x + 2)(x - 2)$
d) $9(x + 2)(x - 2)$
e) $(x + \sqrt{5})(x - \sqrt{5})$
9 a) $\sqrt{6}$
b) 5
c) $\sqrt{5}$
d) $4\sqrt{3}$
e) $8 + 2\sqrt{7}$
10 a) $\frac{3\sqrt{2}}{2}$
b) $\frac{\sqrt{10}}{4}$
c) $\frac{6 - 2\sqrt{6}}{3}$
d) $\frac{\sqrt{2} + \sqrt{10}}{-4}$
11 a) $x = 2$
b) $x = 10$
c) $x = -5$
d) $x = -5$
d) $x = -5$
d) $x = \frac{5y - 2}{3}$
c) $x = \pm \sqrt{\frac{y - 1}{2z}}$
d) $x = \frac{2y + 1}{y - 3}$

b) x = -5 or x = -1c) x = 2.5 or x = -1d) $x = \frac{4}{3}$ or x = 314 a) x = 2.32 or x = -4.32b) x = 2.69 or x = -0.1915 a) $x = 2 + \sqrt{6}$ or $x = 2 - \sqrt{6}$ b) $x = -1 + \frac{3}{\sqrt{2}}$ or $x = -1 - \frac{3}{\sqrt{2}}$ 16 a) $x^2 + 6x + 8 = (x + 3)^2 - 1$ b) -2, 0(-4, 0 (-3, -1) 17 a) $3ab^2$ b) $\frac{1}{8y}$ c) $\frac{x-4}{x-5}$ 18 a) 6*ab* b) $\frac{x}{3}$ c) 27 d) $\frac{5x^2 - x - 3}{x^2(x+1)}$ 19 a) $x \le -1$ b) *x* < 8 c) -3 < x < 3d) $x \le -2$ or $x \ge 2$ e) $x \le 1$ or $x \ge 5$ 20 = 3x - 1 $y = \frac{1}{5}x + 2$ 1. $\frac{1}{3}x$ 0 -2 -1 5 21 a) x = 2, y = -2b) x = 11, y = 16c) x = -3, y = 12 or x = 5, y = 28d) x = 0, y = -2 or x = 3, y = 422 Take three consecutive odd numbers: 2n + 1, 2n + 3 and 2n + 5, where *n* is an integer. 2n + 1 + 2n + 3 + 2n + 5 = 6n + 9 = 3(2n + 3)The sum of three consecutive odd numbers can be written as 3x, where x = 2n + 3. Therefore it is a multiple of 3.

23 E.g. Let x = 3 and y = -1. So $xy = -3 \Rightarrow xy < y$. So Naveen is wrong.



 $31 \quad 3x + 4y - 25 = 0$



62

38 a) See histogram in part c).





Section 1 — Number

Page 6 — Types of Number

- 1 a) **Rational** it's a terminating decimal.
 - b) Rational it's a recurring decimal.
 - c) **Rational** $\sqrt{64} = 8$ which is an integer.
 - d) Irrational 3 is not a square number.
 - e) **Irrational** π is irrational
- 2 a) True any integer can be written as a fraction over 1.
 b) False recurring decimals are rational.

Page 7 — Fractions

1 a)
$$2\frac{2}{3} \times \frac{1}{4} = \frac{8}{3} \times \frac{1}{4} = \frac{2}{3} \times \frac{1}{1} = \frac{2}{3}$$

b) $5\frac{1}{3} \div 2\frac{1}{4} = \frac{16}{3} \div \frac{9}{4} = \frac{16}{3} \times \frac{4}{9} = \frac{64}{27}$
c) $\frac{3}{4} + \frac{1}{3} = \frac{9}{12} + \frac{4}{12} = \frac{13}{12}$
d) $1\frac{1}{6} + 2\frac{1}{2} = \frac{7}{6} + \frac{5}{2} = \frac{7}{6} + \frac{15}{6} = \frac{22}{6} = \frac{11}{3}$
e) $1\frac{3}{7} - \frac{2}{9} = \frac{10}{7} - \frac{2}{9} = \frac{90}{63} - \frac{14}{63} = \frac{76}{63}$
f) $\frac{1}{2} + \frac{2}{3} + \frac{3}{4} = \frac{6}{12} + \frac{8}{12} + \frac{9}{12} = \frac{23}{12}$
g) $(\frac{7}{6} - \frac{1}{7}) \div \frac{1}{2} = (\frac{49}{42} - \frac{6}{42}) \div \frac{1}{2} = \frac{43}{42} \div \frac{1}{2} = \frac{43}{42} \times \frac{2}{1} = \frac{43}{21}$
h) $\frac{3}{2} - \frac{1}{9} + 2\frac{1}{3} = \frac{3}{2} - \frac{1}{9} + \frac{7}{3} = \frac{3}{2} - \frac{1}{9} + \frac{21}{9}$
 $= \frac{3}{2} + \frac{20}{9} = \frac{27}{18} + \frac{40}{18} = \frac{67}{18}$

Section 2 — Basic Algebra

Page 8 — Laws of Indices

1 a) $b^5 \times b^6 = b^{(5+6)} = b^{11}$ b) $a^9 \times a \times b^5 = a^{(9+1)} \times b^5 = a^{10}b^5$ c) $c^5 \div c^2 = c^{(5-2)} = c^3$ d) $9y^{10} \div 3y^{-2} = (9 \div 3) \times y^{(10-(-2))} = 3y^{12}$ e) $x^2 \times x^3 \div x^4 = x^{(2+3-4)} = x$ f) $z^3 \times (y+2)^5 \times z \div (y+2)^2 = z^{(3+1)} \times (y+2)^{(5-2)}$ $= z^4(y+2)^3$ g) $a^{(x+2)} \times a^{2x} = a^{(x+2)+2x} = a^{3x+2}$ h) $x^{-2}y^5 \times x^5y^2 = x^{(-2+5)}y^{(5+2)} = x^3y^7$

Page 9 — Laws of Indices

1 a)
$$a^{0} \div b^{-2} = 1 \div \frac{1}{b^{2}} = b^{2}$$

b) $(4^{x})^{x} = 4^{x^{2}}$
c) $3^{m} \times 9^{2} = 3^{m} \times (3^{2})^{2} = 3^{m} \times 3^{4} = 3^{m+4}$
d) $\left(\frac{1}{a}\right)^{2} \times a^{-3} = a^{-2} \times a^{-3} = a^{-2-3} = a^{-5}$
e) $\left(\frac{1}{z^{9}}\right)^{\frac{1}{3}} = \frac{1^{\frac{1}{3}}}{(z^{9})^{\frac{1}{3}}} = \frac{1}{z^{3}}$
2 a) $4^{\frac{1}{2}} = \sqrt{4} = 2$
b) $27^{\frac{2}{3}} = (\sqrt[3]{27})^{2} = (3)^{2} = 9$
c) $2 \times 32^{\frac{3}{5}} = 2 \times (2^{5})^{\frac{3}{5}} = 2 \times 2^{3} = 2^{4} = 16$
d) $\left(\frac{125}{8}\right)^{\frac{1}{3}} = \sqrt[3]{\frac{125}{8}} = \frac{\sqrt[3]{125}}{\sqrt[3]{8}} = \frac{5}{2}$
e) $\left(\frac{25}{4}\right)^{\frac{-3}{2}} = \left(\frac{4}{25}\right)^{\frac{3}{2}} = \left(\sqrt{\frac{4}{25}}\right)^{3} = \left(\frac{\sqrt{4}}{\sqrt{25}}\right)^{3} = \left(\frac{2}{5}\right)^{3} = \frac{8}{125}$
f) $\left(\frac{16}{9}\right)^{\frac{-3}{2}} = \left(\frac{9}{16}\right)^{\frac{3}{2}} = \left(\sqrt{\frac{9}{16}}\right)^{3} = \left(\frac{\sqrt{9}}{\sqrt{16}}\right)^{3} = \left(\frac{3}{4}\right)^{3} = \frac{27}{64}$
3 $(5^{\frac{1}{4}})^{2} \times (5^{\frac{2}{3}})^{\frac{3}{4}} \div (5^{-1})^{-2} = 5^{\frac{2}{4}} \times 5^{\frac{(2^{3} - 3)}{4}} \div 5^{-1 \times -2}$
 $= 5^{\frac{1}{2}} \times 5^{-\frac{1}{2}} \div 5^{2}$
 $= 5^{0} \div 5^{2} = 5^{-2}$

Page 10 — Multiplying Out Brackets

1 a)
$$(y + 3)(y - 6) = y^2 - 6y + 3y - 18 = y^2 - 3y - 18$$

b) $(a - 3)(b + 4) = ab + 4a - 3b - 12$
c) $(p - 1)(p - 2q)^2 = (p - 1)(p - 2q)(p - 2q)$
 $= (p - 1)(p^2 - 4pq + 4q^2)$
 $= p(p^2 - 4pq + 4q^2) + (-1)(p^2 - 4pq + 4q^2)$
 $= p^3 - 4p^2q + 4pq^2 - p^2 + 4pq - 4q^2$
d) $(s^2 + s + 2)(2s^2 - 2s + 4)$
 $= s^2(2s^2 - 2s + 4) + s(2s^2 - 2s + 4) + 2(2s^2 - 2s + 4)$
 $= (2s^4 - 2s^3 + 4s^2) + (2s^3 - 2s^2 + 4s) + (4s^2 - 4s + 8)$
 $= 2s^4 + 6s^2 + 8$

Page 11 — Factorising

1 a) $20x^2 - 4x = 4x(5x - 1)$ b) $8x^2y + 28xy^2 = 4xy(2x + 7y)$ c) $3\pi a^2 + 4\pi ab + 2\pi a = \pi a(3a + 4b + 2)$ d) $5x^2(x - 1) - 2x(x - 1) = x(x - 1)(5x - 2)$ e) $x^2 - 9 = (x + 3)(x - 3)$ f) $9x^2 - 25 = (3x + 5)(3x - 5)$ g) $p^2 - 49q^2 = p^2 - (7q)^2 = (p + 7q)(p - 7q)$ h) $v^2 - 7u^2 = (v + \sqrt{7}u)(v - \sqrt{7}u)$

Page 12 — Surds

1 a)
$$2\sqrt{24} + 3\sqrt{96} = 2\sqrt{4 \times 6} + 3\sqrt{16 \times 6}$$

 $= 2\sqrt{4}\sqrt{6} + 3\sqrt{16}\sqrt{6}$
 $= 4\sqrt{6} + 12\sqrt{6} = 16\sqrt{6}$
b) $\frac{\sqrt{120}}{\sqrt{15}\sqrt{2}} = \frac{\sqrt{120}}{\sqrt{30}} = \sqrt{\frac{120}{30}} = \sqrt{4} = 2$
c) $(1 + \sqrt{x})^2 - 2\sqrt{x} = (1 + \sqrt{x})(1 + \sqrt{x}) - 2\sqrt{x}$
 $= 1 + 2\sqrt{x} + (\sqrt{x})^2 - 2\sqrt{x} = 1 + x$
2 $\frac{\sqrt{3}}{\sqrt{20}} = \frac{\sqrt{3}}{\sqrt{4}\sqrt{5}} = \frac{\sqrt{3}}{2\sqrt{5}} = \frac{\sqrt{3}\sqrt{5}}{2\sqrt{5}\sqrt{5}} = \frac{\sqrt{15}}{2(\sqrt{5})^2}$
 $= \frac{\sqrt{15}}{2 \times 5} = \frac{\sqrt{15}}{10}$
3 $(\sqrt{7} + \sqrt{12})^2 = (\sqrt{7} + \sqrt{12})(\sqrt{7} + \sqrt{12})$
 $= (\sqrt{7})^2 + 2\sqrt{7}\sqrt{12} + (\sqrt{12})^2$
 $= 7 + 2\sqrt{84} + 12$
 $= 19 + 2\sqrt{4}\sqrt{21}$
 $= 19 + 4\sqrt{21}$

Page 13 — Surds

¹ a)
$$\frac{1}{1-\sqrt{5}} = \frac{1 \times (1+\sqrt{5})}{(1-\sqrt{5})(1+\sqrt{5})}$$

= $\frac{1+\sqrt{5}}{1-(\sqrt{5})^2} = \frac{1+\sqrt{5}}{1-5} = -\frac{1+\sqrt{5}}{4}$

b)
$$\frac{\sqrt{10}}{4 + \sqrt{40}} = \frac{\sqrt{10}}{4 + \sqrt{4}\sqrt{10}} = \frac{\sqrt{10}}{4 + 2\sqrt{10}}$$
$$= \frac{\sqrt{10}(4 - 2\sqrt{10})}{(4 + 2\sqrt{10})(4 - 2\sqrt{10})}$$
$$= \frac{4\sqrt{10} - 20}{4^2 - (2\sqrt{10})^2} = \frac{4\sqrt{10} - 20}{16 - 4 \times 10}$$
$$= \frac{4\sqrt{10} - 20}{-24} = \frac{\sqrt{10} - 5}{-6} = \frac{5 - \sqrt{10}}{6}$$

^{c)}
$$\frac{1+\sqrt{7}}{5+\sqrt{7}} = \frac{(1+\sqrt{7})(5-\sqrt{7})}{(5+\sqrt{7})(5-\sqrt{7})}$$
$$= \frac{5-\sqrt{7}+5\sqrt{7}-(\sqrt{7})^2}{5^2-(\sqrt{7})^2}$$
$$= \frac{5+4\sqrt{7}-7}{25-7} = \frac{-2+4\sqrt{7}}{18} = \frac{2\sqrt{7}-1}{9}$$

d)
$$\frac{2+2\sqrt{2}}{2-2\sqrt{2}} = \frac{(2+2\sqrt{2})(2+2\sqrt{2})}{(2-2\sqrt{2})(2+2\sqrt{2})}$$
$$= \frac{4+4\sqrt{2}+4\sqrt{2}+(2\sqrt{2})^{2}}{4-(2\sqrt{2})^{2}}$$
$$= \frac{4+8\sqrt{2}+8}{4-8} = \frac{12+8\sqrt{2}}{-4} = -3-2\sqrt{2}$$
$$\frac{4+8\sqrt{2}+8}{4-8} = \frac{4}{-4}$$

$$2 \qquad \frac{4}{1-2\sin 60^{\circ}} = \frac{4}{1-2\left(\frac{\sqrt{3}}{2}\right)}$$
$$= \frac{4}{1-\sqrt{3}} = \frac{4(1+\sqrt{3})}{(1-\sqrt{3})(1+\sqrt{3})}$$
$$= \frac{4+4\sqrt{3}}{1-(\sqrt{3})^{2}} = \frac{4+4\sqrt{3}}{1-3}$$
$$= \frac{4+4\sqrt{3}}{-2} = -2-2\sqrt{3}$$

Answers

12

Page 14 — Solving Equations

1 a)
$$4(2x - 3) = 7x \Rightarrow 8x - 12 = 7x \Rightarrow 8x = 7x + 12 \Rightarrow x =$$

b) $3(x + 14) = x + 12 \Rightarrow 3x + 42 = x + 12$
 $\Rightarrow 2x = -30 \Rightarrow x = -15$
c) $\frac{b-7}{3} + \frac{b+1}{5} = -1 \Rightarrow 5(b-7) + 3(b+1) = -15$
 $\Rightarrow 5b - 35 + 3b + 3 = -15$
 $\Rightarrow 8b - 32 = -15 \Rightarrow 8b = 17$
 $\Rightarrow b = \frac{17}{8}$
d) $\frac{q(q+7)}{7} - q = 4 - q^2 \Rightarrow q(q+7) - 7q = 28 - 7q^2$
 $\Rightarrow q^2 + 7q - 7q = 28 - 7q^2$
 $\Rightarrow 8q^2 = 28 \Rightarrow q^2 = \frac{28}{8} = \frac{7}{2}$
 $\Rightarrow 8q^2 = 28 \Rightarrow q^2 = \frac{28}{8} = \frac{7}{2}$
 $\Rightarrow q = \pm \sqrt{\frac{7}{2}}$
2 The area of the whole square will be x^2 .
If the area of $\frac{1}{4}$ of the square is 25, then $\frac{x^2}{4} = 25$
 $\Rightarrow x^2 = 100 \Rightarrow x = \pm 10$.

So the sides are **10 cm**.

Page 15 — Rearranging Formulas

1 a)
$$b(a + 2) = 4 \Rightarrow ab + 2b = 4 \Rightarrow ab = 4 - 2b \Rightarrow a = \frac{4}{b} - 2$$

b) $f = \frac{9}{5}c + 32 \Rightarrow 5f = 9c + 160 \Rightarrow 9c = 5f - 160$
 $\Rightarrow c = \frac{5f - 160}{9}$
2 a) $y = \frac{2x^2 - 3}{x^2 - 1} \Rightarrow y(x^2 - 1) = 2x^2 - 3$
 $\Rightarrow x^2y - y = 2x^2 - 3$
 $\Rightarrow x^2y - 2x^2 = y - 3$
 $\Rightarrow (y - 2)x^2 = y - 3$
 $\Rightarrow x^2 = \frac{y - 3}{y - 2} \Rightarrow x = \pm \sqrt{\frac{y - 3}{y - 2}}$
b) $s = \frac{\sqrt{t + u}}{u} \Rightarrow s^2 = \frac{t + u}{u^2} \Rightarrow s^2u^2 = t + u \Rightarrow t = s^2u^2 - u$

Section 3 — Quadratic Equations

Page 16 — Factorising Quadratics

1 a)
$$x^2 + 3x + 2 = 0 \Rightarrow (x + 1)(x + 2) = 0 \Rightarrow x = -1$$
 and $x = -2$
b) $x^2 + 8x + 7 = 0 \Rightarrow (x + 1)(x + 7) = 0 \Rightarrow x = -1$ and $x = -7$
c) $x^2 = 2(7x - 20) \Rightarrow x^2 - 14x + 40 = 0 \Rightarrow (x - 4)(x - 10) = 0$
 $\Rightarrow x = 4$ and $x = 10$
d) $x^2 - x = 6 \Rightarrow x^2 - x - 6 = 0 \Rightarrow (x + 2)(x - 3) = 0$
 $\Rightarrow x = -2$ and $x = 3$
e) $7x = x^2 + 10 \Rightarrow x^2 - 7x + 10 = 0 \Rightarrow (x - 2)(x - 5) = 0$
 $\Rightarrow x = 2$ and $x = 5$
f) Multiply everything by 2 to get $x^2 + 4x - 12 = 0$.
 $x^2 + 4x - 12 = 0 \Rightarrow (x - 2)(x + 6) = 0 \Rightarrow x = 2$ and $x = -6$
g) Multiply everything by x to get $x^2 - 4x - 12 = 0$.
 $x^2 - 4x - 12 = 0 \Rightarrow (x + 2)(x - 6) = 0 \Rightarrow x = -2$ and $x = 6s$
h) $2x^2 - 2x - 4 = 0 \Rightarrow x^2 - x - 2 = 0$
 $\Rightarrow (x - 2)(x + 1) = 0 \Rightarrow x = 2$ and $x = -1$
2 $x^2 - 2xz + z^2 = (x - z)(x - z) = (x - z)^2$

$$3 \quad -m^2 + 13m - 30 = 0 \Rightarrow m^2 - 13m + 30 = 0$$

$$\Rightarrow (m - 3)(m - 10) \Rightarrow m = 3 \text{ and } 10. \text{ So the temperature}$$

was 0°C after **3 minutes** and after **10 minutes.**

Page 17 — Factorising Quadratics

1 a)
$$2x^2 + 9x + 9 = 0 \Rightarrow (2x + 3)(x + 3) = 0$$

 $\Rightarrow x = -\frac{3}{2} \text{ and } x = -3$
b) $5x^2 + 13x + 6 = 0 \Rightarrow (5x + 3)(x + 2) = 0$
 $\Rightarrow x = -\frac{3}{5} \text{ and } x = -2$
c) $2x^2 = x + 10 \Rightarrow 2x^2 - x - 10 = 0 \Rightarrow (2x - 5)(x + 2)$
 $\Rightarrow x = \frac{5}{2} \text{ and } x = -2$

d)
$$3x + \frac{21}{x} = 16$$
, so multiply through by x:
 $3x^2 + 21 = 16x \Rightarrow 3x^2 - 16x + 21 = 0$
 $\Rightarrow (3x - 7)(x - 3) = 0$
 $\Rightarrow x = \frac{7}{3}$ and $x = 3$.

2 The plane hits the ground when $-4t^2 + 2t + 2 = 0$. Divide through by $-2: 2t^2 - t - 1 = 0$ $\Rightarrow (2t + 1)(t - 1) = 0 \Rightarrow t = 1$ or $t = -\frac{1}{2}$. t can't be negative, so the plane hits the ground **1 second** after being thrown.

Page 18 — The Quadratic Formula

1 a)
$$4x^2 - 7x + 1 = 0 \Rightarrow a = 4, b = -7, c = 1$$

 $x = \frac{-(-7) \pm \sqrt{(-7)^2 - 4 \times 4 \times 1}}{2 \times 4} = \frac{7 \pm \sqrt{49 - 16}}{8} = \frac{7 \pm \sqrt{33}}{8}$
So $x = \frac{7 \pm \sqrt{33}}{8} = 1.593$ (3 d.p.) and
 $x = \frac{7 - \sqrt{33}}{8} = 0.157$ (3 d.p.)

b)
$$6x^2 + x = 4 \Rightarrow 6x^2 + x - 4 = 0 \Rightarrow a = 6, b = 1, c = -4$$

 $x = \frac{-1 \pm \sqrt{1^2 - 4 \times 6 \times -4}}{2 \times 6} = \frac{-1 \pm \sqrt{1 + 96}}{12} = \frac{-1 \pm \sqrt{97}}{12}$
So $x = \frac{\sqrt{97} - 1}{12} = 0.737$ (3 d.p.) and
 $x = \frac{-\sqrt{97} - 1}{12} = -0.904$ (3 d.p.)

c)
$$-2x^2 = 3x - 4 \Rightarrow 2x^2 + 3x - 4 = 0 \Rightarrow a = 2, b = 3, c = -4$$

 $x = \frac{-3 \pm \sqrt{3^2 - 4 \times 2 \times -4}}{2 \times 2} = \frac{-3 \pm \sqrt{9 + 32}}{4} = \frac{-3 \pm \sqrt{41}}{4}$
So $x = \frac{\sqrt{41} - 3}{4} = 0.851$ (3 d.p.) and
 $x = \frac{-\sqrt{41} - 3}{4} = -2.351$ (3 d.p.)

2 $5x^2 + 7x + 3$ has discriminant $7^2 - 4 \times 5 \times 3 = -11$. This is negative, so the quadratic has **no real roots**.

Page 19 — Completing the Square

- 1 a) $x^2 + 4x + 1 \Rightarrow$ Initial bracket is $(x + 2)^2 = x^2 + 4x + 4$ The adjusting number is 1 - 4 = -3 $\Rightarrow x^2 + 4x + 1 = (x + 2)^2 - 3$
 - b) $x^2 12x + 5 \Rightarrow$ Initial bracket is $(x 6)^2 = x^2 12x + 36$ The adjusting number is 5 - 36 = -31 $\Rightarrow x^2 + 12x + 5 = (x - 6)^2 - 31$
 - c) $x^2 20x 10 \Rightarrow$ Initial bracket is $(x 10)^2 = x^2 20x + 100$ The adjusting number is -10 - 100 = -110 $\Rightarrow x^2 - 20x - 10 = (x - 10)^2 - 110$
 - d) $x^2 + 7x 3 \Rightarrow$ Initial bracket is $(x + \frac{7}{2})^2 = x^2 + 7x + \frac{49}{4}$ The adjusting number is $-3 - \frac{49}{4} = -\frac{61}{4}$ $\Rightarrow x^2 + 7x - 3 = (x + \frac{7}{2})^2 - \frac{61}{4}$
- 2 $x^{2} + 6x 8 \Rightarrow$ Initial bracket is $(x + 3)^{2}$ The adjusting number is -8 - 9 = -17 $\Rightarrow x^{2} + 6x - 8 = (x + 3)^{2} - 17$ $(x + 3)^{2} - 17 = 0 \Rightarrow (x + 3)^{2} = 17$ $\Rightarrow x + 3 = \pm \sqrt{17} \Rightarrow x = -3 + \sqrt{17} \text{ and } x = -3 - \sqrt{17}$

Page 20 — Completing the Square

- 1 a) $x^2 + 16x + 3 \Rightarrow$ Initial bracket is $(x + 8)^2$ The adjusting number is 3 - 64 = -61 $\Rightarrow x^2 + 16x + 3 = (x + 8)^2 - 61$ b) If $x^2 + 16x + 3 = 0$, then $(x + 8)^2 - 61 = 0$ $\Rightarrow (x + 8)^2 = 61 \Rightarrow x + 8 = \pm \sqrt{61} \Rightarrow x = -8 + \sqrt{61}$ or $x = -8 - \sqrt{61}$ c) The coefficient of x^2 is positive, so the graph
 - is u-shaped and has a minimum point. This is when $(x + 8)^2 = 0$, so when x = -8. At this point, y = -61, so the minimum point is at (-8, -61). The *x*-intercepts are the solutions to $x^2 + 16x + 3 = 0$ from part b): $(-8 - \sqrt{61}, 0)$ and $(-8 + \sqrt{61}, 0)$ The *y*-intercept is at x = 0: $y = 0^2 + (16 \times 0) + 3 = 3$, so *y*-intercept is (0, 3).



- 2 a) From the sketch, you can see the minimum point is at (-3, 3) so the adjusting number is q = 3. And it tells you that (-3 + p) = 0, so p = 3.
 - b) From part a), you know that $y = (x + 3)^2 + 3$. At the *y*-intercept, x = 0, so $y = 3^2 + 3 = 12$, which means the *y*-intercept is at (0, 12).

Page 21 — Completing the Square

1 a)
$$2x^2 - 3x - 10 = 2(x^2 - \frac{3}{2}x) - 10$$

 \Rightarrow Initial bracket is $2(x - \frac{3}{4})^2 = 2x^2 - 3x + \frac{9}{8}$
The adjusting number is $-10 - \frac{9}{8} = -\frac{89}{8}$
 $\Rightarrow 2x^2 - 3x - 10 = 2(x - \frac{3}{4})^2 - \frac{89}{8}$
b) $-x^2 - 3x - 1 = -(x^2 + 3x) - 1$
 \Rightarrow Initial bracket is $-(x + \frac{3}{2})^2 = -x^2 - 3x - \frac{9}{4}$
The adjusting number is $-1 - (-\frac{9}{4}) = -1 + \frac{9}{4} = \frac{5}{4}$
 $\Rightarrow -x^2 - 3x - 1 = -(x + \frac{3}{2})^2 + \frac{5}{4}$
c) $5x^2 - x - 3 = 5(x^2 - \frac{1}{5}x) - 3$
 \Rightarrow Initial bracket is $5(x - \frac{1}{10})^2 = 5x^2 - x + \frac{1}{20}$
The adjusting number is $-3 - \frac{1}{20} = -\frac{61}{20}$
 $\Rightarrow 5x^2 - x - 3 = 5(x - \frac{1}{10})^2 - \frac{61}{20}$
d) $-\frac{1}{2}x^2 - 3x - 7 = -\frac{1}{2}(x^2 + 6x) - 7$
 \Rightarrow Initial bracket is $-\frac{1}{2}(x + 3)^2 = -\frac{1}{2}x^2 - 3x - \frac{9}{2}$
The adjusting number is $-7 - (-\frac{9}{2}) = -\frac{5}{2}$
 $\Rightarrow -\frac{1}{2}x^2 - 3x - 7 = -\frac{1}{2}(x + 3)^2 - \frac{5}{2}$
2 $x^2 + y^2 - 2x + 6y - 15 = x^2 - 2x + y^2 + 6y - 15 = 0$
 $\Rightarrow (x - 1)^2 - 1 + (y + 3)^2 - 9 - 15 = 0$
 $\Rightarrow (x - 1)^2 + (y + 3)^2 - 25 = 0$
 $\Rightarrow (x - 1)^2 + (y + 3)^2 = 25$

Section 4 — More Algebra

1 a)
$$\frac{3x^2}{7x} = \frac{3x}{7}$$

b) $\frac{8x+16}{2x-4} = \frac{8(x+2)}{2(x-2)} = \frac{4(x+2)}{x-2}$
c) $\frac{x^2-25}{5(x+5)} = \frac{(x+5)(x-5)}{5(x+5)} = \frac{x-5}{5}$
d) $\frac{3x^2+16x-12}{2x^2+13x+6} = \frac{(3x-2)(x+6)}{(2x+1)(x+6)} = \frac{3x-2}{2x+1}$
e) $\frac{x^3+2x^2+x}{x^2-3x-4} = \frac{x(x^2+2x+1)}{(x-4)(x+1)} = \frac{x(x+1)^2}{(x-4)(x+1)} = \frac{x(x+1)}{x-4}$
2 a) $\frac{x+3}{x^2} \times \frac{x}{4} = \frac{x+3}{x} \times \frac{1}{4} = \frac{x+3}{4x}$
b) $\frac{3x+9}{4} \times \frac{x}{3(x+3)} = \frac{3(x+3)}{4} \times \frac{x}{3(x+3)} = \frac{1}{4} \times \frac{x}{1} = \frac{x}{4}$
c) $\frac{10x}{x^2-9} \div \frac{2x+14}{x-3} = \frac{10x}{x^2-9} \times \frac{x-3}{2x+14}$
 $= \frac{10x}{(x-3)(x+3)} \times \frac{x-3}{2(x+7)}$
 $= \frac{5x}{x+3} \times \frac{1}{x+7} = \frac{5x}{(x+3)(x+7)}$
d) $\frac{x^2+5x+6}{9} \div \frac{x^2-4x-21}{6x-42} = \frac{x^2+5x+6}{9} \times \frac{6x-42}{x^2-4x-21}$
 $= \frac{(x+3)(x+2)}{9} \times \frac{6(x-7)}{(x-7)(x+3)}$
 $= \frac{x+2}{3} \times \frac{2}{1} = \frac{2(x+2)}{3}$
Page 23 — Algebraic Fractions

1 a)
$$3 + \frac{2}{x} = \frac{3x}{x} + \frac{2}{x} = \frac{3x+2}{x}$$

b) $\frac{a}{b} - \frac{2a}{3b} = \frac{3a}{3b} - \frac{2a}{3b} = \frac{a}{3b}$
c) $\frac{1}{x+1} - \frac{3}{x} = \frac{x}{x(x+1)} - \frac{3(x+1)}{x(x+1)} = \frac{x-3x-3}{x(x+1)} = \frac{-2x-3}{x(x+1)}$
d) $\frac{4}{x+2} + \frac{3}{x^2} = \frac{4x^2}{x^2(x+2)} + \frac{3(x+2)}{x^2(x+2)} = \frac{4x^2+3x+6}{x^2(x+2)}$
e) $\frac{1}{x+1} - \frac{1}{x(x+1)} + \frac{1}{x} = \frac{x}{x(x+1)} - \frac{1}{x(x+1)} + \frac{x+1}{x(x+1)}$
 $= \frac{x-1+x+1}{x(x+1)} = \frac{2x}{x(x+1)}$
f) $\frac{5x}{7} + \frac{x+3}{2x} = \frac{10x^2}{14x} + \frac{7(x+3)}{14x} = \frac{10x^2+7x+21}{14x}$
g) $\frac{x-1}{x+8} - \frac{3}{x+1} = \frac{(x-1)(x+1)}{(x+8)(x+1)} - \frac{3(x+8)}{(x+8)(x+1)}$
 $= \frac{x^2-1-3x-24}{(x+8)(x+1)} = \frac{x^2-3x-25}{(x+8)(x+1)}$
h) $1 + \frac{4}{x-4} - \frac{2}{x+2}$
 $= \frac{(x-4)(x+2)}{(x-4)(x+2)} + \frac{4(x+2)}{(x-4)(x+2)} - \frac{2(x-4)}{(x-4)(x+2)}$
 $= \frac{x^2-2x-8+4x+8-2x+8}{(x-4)(x+2)} = \frac{x^2+8}{(x-4)(x+2)}$

Page 25 — Inequalities

1 a) $4 - 8x \le -10x - 6 \Rightarrow 2x \le -10 \Rightarrow x \le -5$ b) $-13 \le 2x - 3 < 11 \Rightarrow -10 \le 2x < 14 \Rightarrow -5 \le x < 7$ c) $-11 < 1 - 3x < 7 \Rightarrow -12 < -3x < 6 \Rightarrow -2 < x < 4$



The set of values where the lines overlap satisfy both inequalities, so $-4 < x \le 1$.

- 3 a) $3x^2 8 < 4 \Rightarrow 3x^2 < 12 \Rightarrow x^2 < 4 \Rightarrow -2 < x < 2$
- b) $2x x^2 + 8 > 0 \Rightarrow x^2 2x 8 < 0$ The graph of $y = x^2 - 2x - 8 = (x - 4)(x + 2)$ is a u-shaped curve that crosses the *x*-axis at x = 4 and x = -2, and the curve is below the *x*-axis between these points, so -2 < x < 4.
- c) $6x \ge 2x^2 + 4 \Longrightarrow 2x^2 6x + 4 \le 0$ The graph of $y = 2x^2 - 6x + 4 = 2(x - 1)(x - 2)$ is a u-shaped curve that crosses the *x*-axis at x = 1 and x = 2, and the curve is below the *x*-axis between these points, so $1 \le x \le 2$.

d)
$$-5x^2 + 4x + 2 < 6x - x^2 \Rightarrow 4x^2 + 2x - 2 > 0$$

 $\Rightarrow 2x^2 + x - 1 > 0$
The graph of $y = 2x^2 + x - 1 = (2x - 1)(x + 1)$ is a u-shaped curve that crosses the *x*-axis at $x = -1$ and $x = \frac{1}{2}$. The curve is above the *x*-axis when *x* is less than -1 and when *x* is more than $\frac{1}{2}$, so $x < -1$ or $x > \frac{1}{2}$.

4 $x^2 + 3x + 7 < 3 - 2x \Rightarrow x^2 + 5x + 4 < 0$ The graph of $y = x^2 + 5x + 4 = (x + 4)(x + 1)$ is a u-shaped curve that crosses the *x*-axis at x = -4 and x = -1, and the curve is below the *x*-axis between these points, so -4 < x < -1.

 $2(3-2x) > 14 \Longrightarrow 6-4x > 14 \Longrightarrow 4x < -8 \Longrightarrow x < -2$

So the values of *x* which satisfy both inequalities are -4 < x < -2.

Page 26 — Graphical Inequalities

1 a) Write the inequality as an equation in the form 'y = ': y = 17 - 5x. Draw the graph of y = 17 - 5x. Decide which side of the line is correct by plugging (0, 0) into the inequality: $5x + y > 17 \Rightarrow 0 > 17$ which is false. So the origin is on the wrong side of the line. So shade the region above y = 17 - 5x.



b) Write the inequality as an equation in the form 'y = ': y = 1 - 4x. Draw the graph of y = 1 - 4x. Use the method from part a) to find and shade the region:



c) Write the inequality as an equation in the form y = 2: $y = \frac{1}{2}x - 4$. Draw the graph of y = 17 - 5x.

Use the method from part a) to find and shade the region:



2 Write the second and third inequalities as equations in the form 'y = ': y = 2 - x and y = x - 10.

Draw the graphs of x = 8, y = 2 - x and y = x - 10.

Decide which side of each line is correct by plugging (0, 0) into each inequality: $x < 8 \Rightarrow 0 < 8$ which is true. So the origin is on the correct side of the line. $x + y > 2 \Rightarrow 0 > 2$ which is false. So the origin is on the wrong side of the line. $x - 4 < y + 6 \Rightarrow -4 < 6$ which is true. So the origin is on the correct side of the line. So shade the region to the left of x = 8and above y = 2 - x and y = x - 10.



Page 27 — Graphical Inequalities

1 a) Write each inequality as an equation in the form 'y = ': $y = x^2$ and y = 4x + 2Draw the graphs of $y = x^2$ and y = 4x + 2. $y = x^2$ is u-shaped and intersects the axes at the origin. Find the region by substituting coordinates into each inequality. Use the origin for y = 4x + 2: $8x + 4 < 2y \Rightarrow 4 < 0$ which is false. So the origin is on the wrong side of the line. The origin is on the line $y = x^2$ so use (1, 2) for $y > x^2$: $y > x^2 \Rightarrow 2 > 1$ which is true. So (1, 2) is on the correct side of the line. So label the region above $y = x^2$ and y = 4x + 2.



- b) Write each inequality as an equation in the form 'y = ': $y = -x^2 + 4$ and y = 4 - x
 - Draw the graphs of $y = -x^2 + 4$ and y = 4 x.

 $y = -x^2 + 4$ is n-shaped. To find where it crosses the *x*-axis, solve $-x^2 + 4 = 0$: $-x^2 + 4 = 0 \Rightarrow x^2 - 4 = 0$

 $\Rightarrow (x + 2)(x - 2) = 0$ so the graph crosses the *x*-axis when

x = 2 and x = -2. The *x*-coordinate of the turning point is halfway between 2 and -2, so at x = 0.

It also crosses the y-axis when x = 0, so y = 4.

This also gives the *y*-coordinate of the turning point as 4. Find the region by substituting (0, 0) into each inequality: $-x^2 + 4 \ge y \implies 4 \ge 0$ which is true.

So the origin is on the correct side of the line.

 $y + 10 < 14 - x \Rightarrow 10 < 14$ which is true.

So the origin is on the correct side of the line. So label the region below $y = -x^2 + 4$ and y = 4 - x.



c) Write each inequality as an equation in the form 'y = ': $y = 8 - 2x^2$ and y = 2x + 6Draw the graphs of $y = 8 - 2x^2$ and y = 2x + 6. Use the method from part b) to find and label the region:



d) Write each inequality as an equation in the form 'y = ': $y = -\frac{x^2}{4}$ and $y = x^2 - 4$.

Draw the graphs of $y = -\frac{x^2}{4}$ and $y = x^2 - 4$.

Use the method from part b) to find and label the region:



2 Write each inequality as an equation in the form 'y = ': $y = -2x^2 + 5x + 3$ and $y = x^2 - x - 2$. Draw the graphs of $y = -2x^2 + 5x + 3$ and $y = x^2 - x - 2$.

Use the method from 1b) to find and shade the region:



Page 28 — Simultaneous Equations

1 a) Write the equations in the correct form and label them: (1) 4x - y = 1 and (2) 5x - 2y = 5 $(1) \times 2: 8x - 2y = 2$ (3) $(3) - (2): 3x = -3 \Longrightarrow x = -1$ Sub into (1): $(4 \times -1) - y = 1 \implies -4 - y = 1 \implies y = -5$ Solution: x = -1, y = -5b) Label the equations: (1) 3x + 8y = 25 and (2) 12x - 10y = 16 $(1) \times 4$: 12x + 32y = 100 (3) $(3) - (2): 42y = 84 \Longrightarrow y = 2$ Sub into (1): $3x + (8 \times 2) = 25 \Rightarrow 3x + 16 = 25$ $\Rightarrow 3x = 9 \Rightarrow x = 3$ Solution: x = 3, y = 2c) Label the equations: (1) 3x - 2y = 8 and (2) 7x - 5y = 17 $(1) \times 5: 15x - 10y = 40 (3)$ $(2) \times 2: 14x - 10y = 34$ (4) (3) - (4): x = 6Sub into (1): $(3 \times 6) - 2y = 8 \Rightarrow 18 - 2y = 8$ $\Rightarrow 2y = 10 \Rightarrow y = 5$ Solution: x = 6, y = 5

d) Write the equations in the correct form and label them:

(1) $\frac{1}{2}x + 2y = 12$ and (2) 4x - y = 11(1) $\times 8: 4x + 16y = 96$ (3) (3) $-(2): 17y = 85 \Rightarrow y = 5$ Sub into (1): $\frac{1}{2}x + (2 \times 5) = 12 \Rightarrow \frac{1}{2}x + 10 = 12$ $\Rightarrow \frac{1}{2}x = 2 \Rightarrow x = 4$ Solution: x = 4, y = 5

Page 29 — Simultaneous Equations

1 a) Label the equations: (1) y = 4x + 4 and (2) $y = x^2 + 3x - 8$ Substitute (1) into (2): $4x + 4 = x^2 + 3x - 8$ $x^2 - x - 12 = 0$ (x - 4)(x + 3) = 0 x = 4 or x = -3Substitute x = 4 into (1): $y = (4 \times 4) + 4 = 20$ Substitute x = -3 into (1): $y = (4 \times -3) + 4 = -8$ Solutions: x = 4, y = 20 and x = -3, y = -8 b) Rearrange the linear equation to get y on its own and label the equations: (1) y = 3x - 3 and (2) $y^2 - x^2 = 0$ Substitute (1) into (2): $(3x - 3)^2 - x^2 = 0$ $8x^2 - 18x + 9 = 0$ Use the quadratic formula with a = 8, b = -18 and c = 9: $x = \frac{-(-18) \pm \sqrt{(-18)^2 - 4 \times 8 \times 9}}{2 \times 8} = \frac{18 \pm \sqrt{36}}{16} = \frac{18 \pm 6}{16}$ x = 1.5 or x = 0.75Substitute x = 1.5 into (1): $y = (3 \times 1.5) - 3 = 1.5$ Substitute x = 0.75 into (1): $y = (3 \times 0.75) - 3 = -0.75$ Solutions: *x* = 1.5, *y* = 1.5 and *x* = 0.75, *y* = -0.75 c) Rearrange the linear equation to get yon its own and label the equations: (1) y = -2x - 7 and (2) $x^{2} + xy - 10 = 0$ Substitute (1) into (2): $x^2 + x(-2x - 7) - 10 = 0$ $-x^2 - 7x - 10 = 0$ $x^2 + 7x + 10 = 0$ (x+5)(x+2) = 0x = -5 or x = -2Substitute x = -5 into (1): $y = (-2 \times -5) - 7 = 3$ Substitute x = -2 into (1): $y = (-2 \times -2) - 7 = -3$ Solutions: x = -5, y = 3 and x = -2, y = -3

Page 31 — Proof

1
$$(n + 1)^2 - (n - 1)^2 = n^2 + 2n + 1 - (n^2 - 2n + 1)$$

= $4n$
= $2(2n)$
 $(n + 1)^2 - (n - 1)^2$ can be written as $2x$, where $x = 2n$ is an integer, so it is even.

2 Call the two odd numbers 2a + 1 and 2b + 1. (2a + 1)(2b + 1) = 4ab + 2a + 2b + 1= 2(ab + a + b) + 1

The product of two odd numbers can be written as 2x + 1, where x is an integer and x = ab + a + b, so is odd.

3 Call the two rational numbers *x* and *y*. Rational numbers can be written as fractions with integers on top and bottom, so write $x = \frac{a}{b}$ and $y = \frac{c}{d}$ where *a*, *b*, *c* and *d* are integers, and *b* and *d* are not 0 (because you can't divide by 0).

 $\frac{a}{b} - \frac{c}{d} = \frac{ad}{bd} - \frac{bc}{bd} = \frac{ad - bc}{bd}$. *ad* and *bc* are the product of integers, so are also integers. So ad - bc is an integer. *bd* is also an integer as it is the product of two integers. It is not 0 because neither *b* nor *d* are 0. **The result of subtracting two rational numbers can be**

written as a fraction with integers on the top and bottom, so it is rational.

- 4 Try n = 3: $2^3 + 1 = 8 + 1 = 9$. 9 is not prime, so Phyllis is **wrong**.
- 5 If x = 1 and y = -2, then $\frac{x}{y} = -\frac{1}{2}$ and $\frac{x^2}{y^2} = \frac{1}{4}$. This means $\frac{x}{y} < \frac{x^2}{y^2}$ so the statement is false.

Page 33 — Functions

- 1 $g(2) = \frac{(11 \times 2) + 13}{21 (7 \times 2)} = \frac{35}{7} = 5$
- 2 $f(3) = (5 \times 3^2) + (12 \times 3) 7 = 45 + 36 7 = 74$
- 3 a) $f(2) = (6 \times 2) + 4 = 12 + 4 = 16$

b)
$$g(4) = \frac{28}{4} = 7$$

c) $fg(x) = f\left(\frac{28}{x}\right) = \left(6 \times \frac{28}{x}\right) + 4 = \frac{168}{x} + 4$
d) $gf(x) = g(6x + 4) = \frac{28}{6x + 4} = \frac{14}{3x + 2}$

4 a) $g(0) = (6 \times 0) - 6 = -6$ $\Rightarrow fg(0) = f(-6) = (4 \times (-6)^2) + 3 = 147$ b) $f(0) = (4 \times 0^2) + 3 = 3 \Rightarrow gf(0) = g(3) = (6 \times 3) - 6 = 12$ c) $f(0) = 3 \Rightarrow f^2(0) = ff(0) = f(3) = (4 \times 3^2) + 3 = 39$ d) $g(0) = -6 \Rightarrow g^2(0) = gg(0) = g(-6) = (6 \times -6) - 6 = -42$

5 a)
$$x = 9 - 11y \Rightarrow y = \frac{9 - x}{11}$$
 so $f^{-1}(x) = \frac{9 - x}{11}$
b) $x = \frac{y - 4}{2} \Rightarrow y = 2x + 4$ so $g^{-1}(x) = 2x + 4$

Section 5 — Graphs

Page 34 — Straight Lines

- 1 Rearrange the equation of the line into y = mx + c form to find the gradient: $2x + y = -2 \Rightarrow y = -2x - 2$, so the gradient is **-2**. When y = 0, $2x = -2 \Rightarrow x = -1$, so *x*-intercept is (**-1**, **0**). When x = 0, y = -2, so *y*-intercept is (**0**, **-2**).
- 2 m = 3, c = 5, so y = 3x + 5.
- 3 $(x_1, y_1) = (3, 4), (x_2, y_2) = (-2, 1).$ So the gradient is $\frac{1-4}{-2-3} = \frac{-3}{-5} = \frac{3}{5}$

Page 35 — Straight Lines

1
$$(x_1, y_1) = (1, 2), (x_2, y_2) = (5, 9).$$

The gradient is $\frac{9-2}{5-1} = \frac{7}{4} \Rightarrow y = \frac{7}{4}x + c$
Plugging in the point (1, 2) gives:
 $2 = \frac{7}{4} + c \Rightarrow c = 2 - \frac{7}{4} = \frac{1}{4}, \text{ so } y = \frac{7}{4}x + \frac{1}{4}.$
2 a) $(x_1, y_1) = (-3, 4), (x_2, y_2) = (7, -2).$
The gradient is $\frac{-2-4}{7-(-3)} = \frac{-6}{10} = -\frac{3}{5} \Rightarrow y = -\frac{3}{5}x + c$
Plugging in the point (-3, 4) gives:
 $4 = -\frac{3}{5} \times -3 + c \Rightarrow 4 = \frac{9}{5} + c \Rightarrow c = 4 - \frac{9}{5} = \frac{11}{5}$
So $y = -\frac{3}{5}x + \frac{11}{5} \Rightarrow 5y = -3x + 11$
 $\Rightarrow 3x + 5y - 11 = 0$
b) $\sqrt{(7-(-3))^2 + (-2-4)^2} = \sqrt{10^2 + (-6)^2} = \sqrt{136}$
 $= 11.66$ to 2 d.p.

3 a) At t = 2, d = 10, so $(t_1, d_1) = (2, 10)$. At t = 3.5, d = 17.5, so $(t_2, d_2) = (3.5, 17.5)$. Gradient of line through these two points $= \frac{d_2 - d_1}{t_2 - t_1} = \frac{17.5 - 10}{3.5 - 2} = \frac{7.5}{1.5} = 5$ So the equation so far is d = 5t + c. Substitute (2, 10) into the equation and solve for c: $d = 5t + c \Rightarrow 10 = 5 \times 2 + c \Rightarrow c = 10 - 10 = 0$. So the equation of the line is d = 5t.

b) $12.5 = 5t \Rightarrow t = 2.5$ hours

Page 36 — Parallel and Perpendicular Lines

 $5y + 3x - 7 = 0 \Rightarrow y = -\frac{3}{5}x + \frac{7}{5}$ 1 Line F's gradient is $-\frac{3}{5}$, so line G's gradient is $-\frac{3}{5}$. Line G goes through the origin, so c = 0. The equation of line G is $y = -\frac{3}{5}x$. Line P is given by y = 4x + c. Plug in point (-1, -2): 2 $-2 = -4 + c \Rightarrow c = 2$, so line P's equation is y = 4x + 2. Rearrange to get: 4x - y + 2 = 0. Line Q has gradient $-1 \div 4 = -\frac{1}{4}$, so line Q's equation is y $= -\frac{1}{4}x + c$. Plugging in the point (-1, -2) gives: $-2 = \frac{1}{4} + c \Rightarrow c = -\frac{9}{4}$. So line Q's equation is $y = -\frac{1}{4}x - \frac{9}{4}$. Rearrange to get: x + 4y + 9 = 0. 3 $(x_1,\,y_1)=(4,\,3),\,(x_2,\,y_2)=(9,\,5)$ l_1 's gradient is $\frac{5-3}{9-4} = \frac{2}{5}$, so l_2 's gradient is $-1 \div \frac{2}{5} = -1 \times \frac{5}{2} = -\frac{5}{2}$ l_2 is given by $y = -\frac{5}{2}x + c$. Plug in point Q (9, 5): $5 = -\frac{45}{2} + c \Rightarrow c = 5 + \frac{45}{2} = \frac{55}{2},$ so *l*₂ is given by $y = -\frac{5}{2}x + \frac{55}{2}$.

Rearrange to get: 5x + 2y - 55 = 0.

Page 37 — Quadratic Graphs

- 1 a) The x^2 coefficient is positive, so this is a u-shaped quadratic with a minimum point. $x = 0 \Rightarrow y = 0^2 - 3 \times 0 = 0 \Rightarrow y$ -intercept is at (0, 0).
 - Solve by factorising to find the *x*-intercept is at (0, 0). Solve by factorising to find the *x*-intercepts: $x^2 - 3x = x(x - 3) = 0 \Rightarrow x$ -intercepts at (0, 0) and (3, 0). The minimum point is half way between the *x*-intercepts, so is at $x = \frac{3+0}{2} = \frac{3}{2}$. $x = \frac{3}{2} \Rightarrow y = \left(\frac{3}{2}\right)^2 - 3 \times \frac{3}{2} = \frac{9}{4} - \frac{9}{2} = -\frac{9}{4}$,

so the minimum point is at
$$(\frac{3}{2}, -\frac{9}{4})$$
 [or (1.5, -2.25)].



b) Use the method from part a) to sketch and label the graph.



c) Use the method from part a) to sketch and label the graph. The coefficient of x^2 is negative, so the graph is n-shaped. Also, the quadratic has a double root at x = 3, so this is the vertex and the graph just touches the *x*-axis here.



d) Use the method from part a) to sketch and label the graph.



Page 38 — Harder Graphs

- 1 a) y = (x + 1)(x 1)(x + 2), so the *x*-intercepts are at x = -1, 1 and -2.
 - $x = 0 \Rightarrow y = -2$, so the *y*-intercept is at -2.



b) Use the method from part a) to sketch and label the graph.



c) Factorise the cubic:

 $-x^3 - 5x^2 + 6x = -x(x^2 + 5x - 6) = -x(x + 6)(x - 1)$. Now use the method from part a) to sketch and label the graph. This is a negative cubic, so goes from top left to bottom right.



Page 39 — Harder Graphs

- 1 a) $a = 1, b = 3, r^2 = 16$. So the circle has centre (1, 3) and radius 4.
 - b) a = 0, b = -2, $r^2 = 50$ So the circle has centre (0, -2) and radius $\sqrt{50} = \sqrt{25}\sqrt{2} = 5\sqrt{2}$
 - c) $a = -\frac{1}{2}$, $b = \frac{1}{2}$ and $r^2 = 2$. So the circle has centre $(-\frac{1}{2}, \frac{1}{2})$ and radius $\sqrt{2}$.
- 2 The centre of this circle is at (1, 2). The radius of the circle goes from (1, 2) to (3, 4), so it has gradient $\frac{4-2}{3-1} = \frac{2}{2} = 1$. This means that the tangent at (3, 4) has gradient -1. So the equation of the tangent so far is y = -x + c. Plug in point (3, 4) to get $4 = -3 + c \Rightarrow c = 7$, so the equation of the tangent is y = -x + 7.

Page 40 — Graph Transformations



b) f(x) + 3 is described by the column vector $\begin{pmatrix} 0\\3 \end{pmatrix}$. f(x-2) is described by the column vector $\begin{pmatrix} 2\\0 \end{pmatrix}$.

Page 41 — Graph Transformations

1 The graph of y = -f(x) looks like this:



The graph of y = f(-x) looks like this:



- 2 a) g(x) = g(-x) means that the graph is symmetrical about the *y*-axis. $x^2 4$ could be g(x), because $y = x^2 4$ is symmetrical about the *y*-axis.
 - b) This cannot be g(x) it is not symmetrical about the *y*-axis.
 - c) This **could** be g(x), because $-x^2$ is symmetrical about the *y*-axis.
- 3 a) This is a reflection in the *x*-axis, so the *x*-coordinate stays the same, but the *y*-coordinate is multiplied by -1. So the turning point is at (4, 7).
 - b) This is a reflection in the *y*-axis, so the *y*-coordinate stays the same, but the *x*-coordinate is multiplied by -1. So the turning point is at (-4, -7).

Section 6 — Trigonometry and Vectors

Page 42 — Trigonometry — Sin, Cos, Tan

1 a)
$$\cos 70^\circ = \frac{l}{12} \Rightarrow l = 12 \times \cos 70^\circ = 4.10 \text{ cm (3 s.f.)}$$

b) $\tan 62^\circ = \frac{l}{2} \Rightarrow l = 2 \times \tan 62^\circ = 3.76 \text{ cm (3 s.f.)}$
c) $\sin 49^\circ = \frac{l}{1.5} \Rightarrow l = 1.5 \times \sin 49^\circ = 1.13 \text{ cm (3 s.f.)}$
d) $\cos 37^\circ = \frac{4}{l} \Rightarrow l = \frac{4}{\cos 37^\circ} = 5.01 \text{ cm (3 s.f.)}$
2 $\sin x = \frac{11}{26} \Rightarrow x = \sin^{-1}(\frac{11}{26}) = 25.03^\circ$

Page 43 — Trigonometry — Sin, Cos, Tan

1 cos 170° = -0.9848... , sin 170° = 0.1736... So **Q** = (-0.98, 0.17) (2 d.p.)

Page 44 — Trigonometry — Graphs

1
1

$$y = \sin x$$
 $y = \frac{1}{\sqrt{2}}$
 $y = -0.2$
 $y = -0.7$
 $y = -0.7$

- a) $\sin 45^\circ = \frac{1}{\sqrt{2}}$, so 45° is one solution.
 - Using the sketch, you can see that the other solution is at: $180^{\circ} 45^{\circ} = 135^{\circ}$
- So the solutions are $x = 45^{\circ}$ and $x = 135^{\circ}$.
- b) $\sin^{-1}(-0.7) = -44.42...^{\circ}$. This is outside the interval so add 360° to get the first solution in the interval: -44.42...° + $360 = 315.57...^{\circ}$

Using the sketch above, you can see that the other solution is the same distance to the right of 180° as the first solution is to the left of 360°.

The first solution is $360 - 315.57...^{\circ} = 44.42...^{\circ}$ to the left of 360, so the other solution is $44.42...^{\circ}$ to the right of $180^{\circ}: 180^{\circ} + 44.42...^{\circ} = 224.42...^{\circ}$

- So solutions to 1 d.p. are $x = 315.6^{\circ}$ and $x = 224.4^{\circ}$.
- c) $\sin^{-1}(-0.2) = -11.53...^{\circ}$. This is outside the interval, so add 360° to get the first solution in the interval:

 $-11.53...^{\circ} + 360 = 348.46...^{\circ}$. Using the sketch above, you can see that the other solution is the same distance to the right of 180° as the first solution is to the left of 360° .

The first solution is $360 - 348.46...^{\circ} = 11.53...^{\circ}$ to the left of 360° , so the other solution is $11.53...^{\circ}$ to the right of $180^{\circ}: 180^{\circ} + 11.53...^{\circ} = 191.53...^{\circ}$ So solutions to 1 d.p. are $x = 348.5^{\circ}$ and $x = 191.5^{\circ}$.

Page 45 — Trigonometry — Graphs



- a) $\cos 45^\circ = \frac{1}{\sqrt{2}}$, so 45° is one solution. Using the sketch, you can see that there is another solution at: $360^\circ - 45^\circ = 315^\circ$ There are also solutions at -45° and -315° because the graph is symmetrical about the *y*-axis. So the solutions are $x = 45^\circ$, $x = -45^\circ$, $x = 315^\circ$ and $x = -315^\circ$.
- b) $\cos^{-1}(0.1) = 84.26...^{\circ}$, which gives one solution. Using the sketch above, you can see that there is another solution at $360^{\circ} - 84.26...^{\circ} = 275.73...^{\circ}$. There also solutions at $-84.26...^{\circ}$ and $-275.73...^{\circ}$ because the graph is symmetrical about the *y*-axis. So solutions to 1 d.p. are $x = 84.3^{\circ}$, $x = -84.3^{\circ}$, $x = 275.7^{\circ}$ and $x = -275.7^{\circ}$.
- c) $\cos^{-1}(-0.4) = 113.57...^{\circ}$, which gives one solution. Using the sketch above, you can see that there is another solution at: $360^{\circ} - 113.57...^{\circ} = 246.42...^{\circ}$ There are also solutions at $-113.57...^{\circ}$ and $-246.42...^{\circ}$ because the graph is symmetrical about the *y*-axis. So solutions to 1 d.p. are $x = 113.6^{\circ}$, $x = -113.6^{\circ}$, $x = 246.4^{\circ}$ and $x = -246.4^{\circ}$.

Page 46 — Trigonometry — Graphs



- a) tan⁻¹(-1) = -45° is one solution. Using the sketch, you can see that there are other solutions at: -180° - 45° = -225° 180° - 45° = 135° 360° - 45° = 315°. So the solutions are x = -45°, x = -225°, x = 135° and x = 315°.
 b) tan⁻¹(7) = 81.86...°, which gives one solution. Using the sketch above, you can see that there are other solutions at: -180° + 81.86...° = -98.13...°
 - solutions at: $-180^{\circ} + 81.86...^{\circ} = -98.13...^{\circ}$ $-360^{\circ} + 81.86...^{\circ} = -278.13...^{\circ}$ $180^{\circ} + 81.86...^{\circ} = 261.86...^{\circ}.$ So solutions to 1 d.p. are $x = 81.9^{\circ}$, $x = -98.1^{\circ}$, $x = -278.1^{\circ}$ and $x = 261.9^{\circ}.$
- c) $\tan^{-1}(-6) = -80.53...^{\circ}$, which gives one solution. Using the sketch above, you can see that there are other solutions at: $180^{\circ} - 80.53...^{\circ} = 99.46...^{\circ}$ $360^{\circ} - 80.53...^{\circ} = 279.46...^{\circ}$ $-180^{\circ} - 80.53...^{\circ} = -260.53...^{\circ}$ So solutions to 1 d.p. are $x = -80.5^{\circ}$, $x = 99.5^{\circ}$, $x = 279.5^{\circ}$ and $x = -260.5^{\circ}$.

Page 47 — The Sine and Cosine Rules

- 1 a) Using the sine rule $\frac{x}{\sin 42^\circ} = \frac{12}{\sin 33^\circ}$ $\Rightarrow x = \frac{12 \times \sin 42^\circ}{\sin 33^\circ} = 14.7 \text{ cm (1 d.p.)}$
 - b) The angle opposite x is $180^\circ 23^\circ 101^\circ = 56^\circ$. Using the sine rule gives $\frac{x}{\sin 56^\circ} = \frac{3.5}{\sin 101^\circ}$ $\Rightarrow x = \frac{3.5 \times \sin 56^\circ}{\sin 101^\circ} = 3.0 \text{ m (1 d.p.)}$
 - c) Use the sine rule to find the angle that is opposite the 31 cm side (call it 'A'): $\frac{31}{\sin A} = \frac{46}{\sin 107^{\circ}}$ $\Rightarrow \sin A = \frac{31 \times \sin 107^{\circ}}{46} \Rightarrow A = \sin^{-1} \left(\frac{31 \times \sin 107^{\circ}}{46}\right)$

A = 40.12...°

So the angle opposite *x* is $180^{\circ} - 107^{\circ} - 40.12...^{\circ} = 32.87...^{\circ}$ Use the sine rule again: $\frac{x}{\sin 32.87...^{\circ}} = \frac{46}{\sin 107^{\circ}}$ $\Rightarrow x = \frac{46 \times \sin 32.87...^{\circ}}{\sin 107^{\circ}} = 26.1 \text{ cm (1 d.p.)}$

Page 48 — The Sine and Cosine Rules

1 a) The cosine rule gives $x^2 = 15^2 + 16^2 - 2 \times 15 \times 16 \times \cos 19^\circ$ $\Rightarrow x = \sqrt{27.15...} = 5.2 \text{ mm} (1 \text{ d.p.})$ The area of the triangle is: $\frac{1}{2} \times 15 \times 16 \times \sin 19^\circ = 39.1 \text{ mm}^2$ (1 d.p.) b) Rearrange the cosine rule into the correct form to find an angle: $\cos A = \frac{b^2 + c^2 - a^2}{24}$

to find an angle: $\cos A = \frac{b^2 + c^2 - a^2}{2bc}$ This gives $\cos x = \frac{7^2 + 3^2 - 5^2}{2 \times 7 \times 3} = 0.7857...$

⇒ $x = \cos^{-1} (0.7857...) = 38.21...° = 38.2° (1 d.p.)$ The area of the triangle is:

$$\frac{1}{2} \times 3 \times 7 \times \sin 38.21...^{\circ} = 6.5 \text{ cm}^2$$
 (1 d.p.)

c) The cosine rule gives: $\cos x = \frac{11^{2} + 12^{2} - 13^{2}}{2 \times 11 \times 12} = 0.363...$ $\Rightarrow x = \cos^{-1} (0.363...) = 68.67...^{\circ} = 68.7^{\circ} (1 \text{ d.p.})$ The area of the triangle is: $\frac{1}{2} \times 12 \times 11 \times \sin 68.67...^{\circ} = 61.5 \text{ cm}^{2} (1 \text{ d.p.})$

Page 49 — Vectors

- 1 $\overrightarrow{AC} = \overrightarrow{AB} + \overrightarrow{BC} = 2\mathbf{s} + \mathbf{t} + 2\mathbf{t} \frac{1}{2}\mathbf{s} = \frac{3}{2}\mathbf{s} + 3\mathbf{t}.$ So $2\overrightarrow{AC} = 3\mathbf{s} + 6\mathbf{t} = 3(\mathbf{s} + 2\mathbf{t})$ as required.
- 2 $\overrightarrow{\mathsf{TW}} = \overrightarrow{\mathsf{TU}} + \overrightarrow{\mathsf{UV}} + \overrightarrow{\mathsf{VW}} = \mathbf{a} + \mathbf{b} (-2\mathbf{a})$ = $\mathbf{a} + 2\mathbf{a} + \mathbf{b} = 3\mathbf{a} + \mathbf{b}$

Page 50 — Vectors

- 1 $\mathbf{v} = 4\mathbf{a} + 6\mathbf{b}$, $\mathbf{u} = 6\mathbf{a} + 9\mathbf{b}$. $\mathbf{v} = 2(2\mathbf{a} + 3\mathbf{b})$, $\mathbf{u} = 3(2\mathbf{a} + 3\mathbf{b})$ so $\mathbf{v} = \frac{2}{3}\mathbf{u}$. \mathbf{v} and \mathbf{u} are scalar multiples of each other, which means they are parallel.
- 2 To show the W, X and Y are collinear, you need to show \overrightarrow{YX} is parallel to \overrightarrow{XW} . $\overrightarrow{YX} = \overrightarrow{YO} + \overrightarrow{OX} = -2\mathbf{v} + \mathbf{u} - \mathbf{v} = \mathbf{u} - 3\mathbf{v}$ $\overrightarrow{XW} = \overrightarrow{XO} + \overrightarrow{OW} = -(\mathbf{u} - \mathbf{v}) + 2\mathbf{u} - 4\mathbf{v} = \mathbf{u} - 3\mathbf{v}$ This shows that \overrightarrow{YX} is parallel to \overrightarrow{XW} and **W**, X and Y are collinear.

3 a)
$$\sqrt{3^2 + 2^2} = \sqrt{13}$$

b) $\sqrt{(-1)^2 + (-1)^2} = \sqrt{2}$

Section 7 — Statistics and Probability

Page 51 — Sampling

E.g. This sample may be biased because it hasn't been chosen at random from residents of the town. / The sampling method excludes residents who don't know about the meeting, or who are unable to attend, e.g. because they are at work. Attendees are therefore more likely to be older or unemployed. / It's possible that people who live outside the town may also attend the meeting, which would lead to sampling from the wrong population. / It's likely to be biased towards people with strong views, as they are more likely to choose to attend the meeting. / If only a small number of people attend the meeting, the sample will be too small to represent the population.

Page 52 — Data Basics

210-320

1

Lower Upper Length Class Class Frequency B'darv B'darv (mm)(mm) (mm) 80-110 15 75 115 120-200 16 115 205

205

325

19

Class

Mid-

point

(mm)

95

160

265

Class

Width

(mm)

40

90

120

Page 53 — Histograms



74

Height (<i>h</i> , cm)	Frequency	Frequency density
$150 < h \le 170$	40	$40 \div 20 = 2$
$170 < h \le 190$	32	$32 \div 20 = 1.6$
$190 < h \le 210$	26	26 ÷ 20 = 1.3
$210 < h \le 240$	9	$9 \div 30 = 0.3$





Between 155 cm and 170 cm: $2 \times (170 - 155) = 2 \times 15 = 30$ All of the 170 < $h \le 190$ class is included, which is another 32 people. Between 190 cm and 200 cm: $1.3 \times (200 - 190) = 1.3 \times 10 = 13$. So the estimate is 30 + 32 + 13 = 75 people.

Page 54 — Averages

- 1 Put the data into ascending order: 1, 4, 5, 5, 6, 7, 8 The **mode** is **5**. There are 7 data values, so the middle value is the fourth one, so the **median** is **5**. Calculate the **mean**: $(1 + 4 + 5 + 5 + 6 + 7 + 8) \div 7 = 5.14$ (2 d.p.)
- 2 Plug the values into the formula: $\overline{x_1} = 12, n_1 = 5, \ \overline{x_2} = 15, n_2 = 6$ $\overline{x} = \frac{5 \times 12 + 6 \times 15}{5 + 6} = \frac{150}{11} = 13.64$ (2 d.p.)

Page 55 — Averages

1 a) The total of the frequencies is 30, so the median is halfway between the 15th and 16th values. This means the median is in the **50-90 g class**.

1 .				
b)	Mass (g)	Frequency f	mid-point x	fx
	0-40	6	$(45 + 0) \div 2 = 22.5$	135
	50-90	11	$(45 + 95) \div 2 = 70$	770
	100-140	8	$(95 + 145) \div 2 = 120$	960
	150-190	5	$(145 + 195) \div 2 = 170$	850
	Total	30	_	2715

So $\overline{x} = 2715 \div 30 = 90.5 \text{ g}$

Answers

Page 56 — Cumulative Frequency





b) The median is half way up the side of the graph, so read along from 30 to get approximately 450 mm.
The upper quartile is 75% of the way up the side of the graph, so read along from 45 to get approximately 482 mm.
The lower quartile is 25% of the way up the side of

the graph, so read along from 15 to get approximately 405 mm.

So the interquartile range is approximately 482 - 405 = 77 mm.

Page 57 — Probability

- 1 a) There are 2 outcomes where the event 'roll 1 or 2' happens, and 6 outcomes in total. So P(roll 1 or 2) = $\frac{2}{6} = \frac{1}{2}$.
 - So P(roll 1 or 2) = $\frac{2}{6} = \frac{1}{3}$. b) There are 4 outcomes matching 'roll higher than 2', and 6 outcomes in total. So P(roll higher than 2) = $\frac{4}{6} = \frac{2}{3}$.

(Or you could use the fact that probabilities add up to 1. The result of the dice roll will either be a number higher than 2, or it will be equal to 1 or 2, so P(roll higher than 2) = 1 - P(roll 1 or 2)

$$(\operatorname{nan} 2) = 1 - P(\operatorname{roll} 1 \text{ or })$$

= $1 - \frac{1}{3} = \frac{2}{3}$.)

2 There are 4 outcomes where the event 'ball is yellow and labelled A' happens, and 30 outcomes in total. So P(ball is yellow and labelled A) = $\frac{4}{30} = \frac{2}{15}$.

Page 58 — Probability

1 You need two circles — one to show the people who bought fiction books and one to show the people who bought travel books. 2 people bought both types, so 2 goes in the overlap. 22 - 2 = 20 people bought only fiction books and 7 - 2 = 5 people bought only travel books. 40 - 20 - 2 - 5 = 13 people bought neither type of book. So the Venn diagram looks like this:



2 The event 'likes biology or chemistry or both' is shown by the area that's in B or in C or in both circles. Add up the numbers to get: P(likes biology or chemistry or both) = 0.25 + 0.05 + 0.25 + 0.1 + 0.1 + 0.15 = 0.9

Page 59 — Laws of Probability

- 1 If M = 'Martha plays' and S = 'Sahil plays',then P(M or S) = P(M) + P(S) - P(M and S).The events M and S are independent, so $P(M \text{ and } S) = P(M) \times P(S) = 0.3 \times 0.4 = 0.12.$ So P(M or S) = 0.3 + 0.4 - 0.12 = 0.58(Or you could do 1 - P(neither play) to get: $1 - (0.7 \times 0.6) = 1 - 0.42 = 0.58$)
- 2 You need to show that $P(A) \times P(B) = P(A \text{ and } B)$. $P(A) \times P(B) = 0.6 \times 0.5 = 0.3 = P(A \text{ and } B)$, so the events are independent.

Page 60 — Tree Diagrams

1 Draw a set of branches for the first Friday. At the end of each branch, draw a set for the second Friday. Claire has curry this week, so you get the following probabilities:



Now you can multiply along the branches P and P to find P(pizza and pizza) = $0.6 \times 0.25 = 0.15$.

Index

Α

addition law (probability) 59 algebra 8-33 algebraic fractions 22, 23 area of a triangle 48 asymptotes 39 averages 54, 55

В

bias 51 bimodal 54

С

census 51 circle equations 21, 39 class mid-point 52 class width 52, 53 collinear points 50 column vectors 40, 49, 50 common denominators 7, 23 common factors 11, 22 completing the square 19-21 composite functions 32 continuous variables 52 cosine rule 47, 48 cumulative frequency 56

D

denominators 7, 12, 22 dependent events 60 difference of two squares 11, 12 discrete variables 52 discriminants 18 disproof by counter example 31

Ε

events 57 dependent 60 independent 59 mutually exclusive 59

F

factorising 11 quadratics 16, 17 FOIL method 10 fractions 7 algebraic 22, 23 frequency density 53 frequency tables 52, 55 functions 32, 33

G

gradients 34-36 graphical inequalities 26, 27 graphs 34-39 circles 21, 39 cubics 38 exponentials 38 quadratics 20, 25, 27, 37 reciprocals 39 straight lines 26, 34-36 trig 44-46 graph transformations 40, 41 grouped frequency tables 52, 55

Η

histograms 53 hypotenuse 42

imaginary numbers 6 improper fractions 7 independent events 59 indices 8, 9 inequalities 24-27 integers 6, 30 interquartile range 56 inverse functions 33 irrational numbers 6

L

laws of indices 8, 9 lower class boundaries 52 lower quartiles 56

Μ

magnitude of a vector 50 mean 54, 55 median 54-56 mixed numbers 7 modal class 55 mode 54 multiplying out brackets 10 mutually exclusive events 59

Ν

numerators 7, 22

0

outcomes 57

Ρ

parallel lines 36 parallel vectors 50 perpendicular lines 36 populations 51 powers 8, 9 probability 57-60 product law (probability) 59 proof 30, 31 Pythagoras' theorem 35

Q

quadratics 16-21 completing the square 19-21 graphs 20, 25, 27, 37 inequalities 26, 27 quadratic formula 18 sketching 20, 25, 27, 37 qualitative and quantitative variables 52

R

rationalising the denominator 12, 13 rational numbers 6 real numbers 6 rearranging formulas 15 resultant vectors 49

S

sample space 57 sampling 51 scalars 49, 50 simple random sampling 51 simultaneous equations 28, 29 sine rule 47 SOH CAH TOA 42 solving equations 14 surds 12, 13

Т

transformations 40, 41 tree diagrams 60 trigonometry 42-48 graphs 44-46 two-way tables 57

U

unit circle 43 upper class boundaries 52 upper quartiles 56

V

vectors 49, 50 Venn diagrams 58

Υ

y = mx + c 34-36



CGP books — they might just save your life...



OK, maybe not your *life*. But when it comes to exam revision, CGP are the undisputed champions. You can order any of our books (with next-day delivery!) from us, online or by phone:

www.cgpbooks.co.uk • 0800 1712 712

Or you'll find our range in any good bookshop, including:





Waterstones



Jumpstart your A-Level Maths prep with CGP!

You'll be out of the blocks in no time with this brilliant Head Start book...

- Super-handy roundups of essential GCSE skills... Including a quick test to check how much you remember
- Tantalising teasers of new A-Level concepts... It's true — get a first look at how to use them
- Straightforward notes and practice questions... Yep, we'll give those brain cells a real workout

This isn't just any old A-Level Maths book, this is a CGP A-Level Maths book! 🙂

P.S. You're on a roll — why not get a Head Start on **A-Level Science** too with CGP!



CAUTION: Book exterior may be slippery when wet



www.cgpbooks.co.uk